

Antioxidant Potential and Antitumour Activities of *Nendran* Banana Peels in Breast Cancer Cell Line

P. S. KUMAR* S. DURGADEVI¹, A. SARAVANAN AND S. UMA

ICAR-National Research Centre for Banana, Trichy-620 102, ¹Dr. N. G. P. Arts and Science College, Coimbatore-620 148, India

Kumar *et al.*: Anticancer activity of Banana peel

The present study was aimed to evaluate the antioxidant and antitumor activities of the aqueous methanol extract of *Nendran* banana peel as a valuable source for bioactive compounds against cancer. Free radical scavenging activity of the peel extracts was evaluated using 1,1-diphenyl-2-picrylhydrazyl and ferric reducing antioxidant power assay. The antitumor activity on MCF-7 breast cancer cell line was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide and ethidium bromide/acridine orange staining assay. Extract of banana peel was more efficient in reducing ferric ion with the increase in concentration and exhibited IC₅₀ 180.33 µg/ml, free radical inhibition activity under 2,2-diphenyl-1-picrylhydrazyl hydrate assay. Flavonoid content was high (51.47 µg/ml) in the peel extract, which could have contributed to the antioxidant activity. The antioxidant enzyme concentration such as ascorbate peroxidase, catalase, peroxidase and superoxide dismutase were found higher in the fruit pulp than peel. As the concentration of extract increased from 20 to 200 µg/ml, the cell viability of MCF-7 decreased from 91.14 to 24.7 %. The IC₅₀ values of plant extract-treated cells (120.6 µg/ml) showed typical apoptotic and necrotic morphological features such as condensed nuclei, membrane blebbing and formation of apoptotic bodies. This indicated the efficiency of sample against breast cancer with the highest inhibition of growth.

Key words: *Nendran* banana peel, bioactive compounds, apoptotic index, phenols, flavonoids, antioxidant

Cancer is a hyper-proliferative disorder, which involves transformation, deregulation of apoptosis, proliferation, invasion, angiogenesis, and metastasis. Worldwide, more than half a million people died of cancer every year, at a rate of 1500 patients every day^[1]. Breast cancer is the second leading cancer in women, accounting for 25 % of all cases^[2]. Currently tamoxifen, selective oestrogen receptor modulator, is widely used in the prevention and treatment of oestrogen receptor positive breast cancer^[3]. However, a significant number of patients develop tamoxifen resistance and experience severe side effects^[4,5]. It is therefore, imperative to search for newer compounds, which are effective, greener and not with many side effects.

As the conventional cancer therapies failed to fulfil the criteria for successful cancer treatments, recent research revolved around the urgency to develop suitable therapy for the cure of cancer with no toxic effects through methodical and scientific exploration of the enormous pool of synthetic, biological, and natural products^[2,6-9]. Plant-derived natural products

received significant attention due to their diverse pharmacological properties and cancer preventive effects. With the presence of flavonoids, alkaloids and terpenes, these offer different forms of cytotoxic effects such as cell proliferation inhibition, apoptosis induction or modulating signal transduction, without exhibiting considerable damage to normal cells^[7-13]. These are relatively new, but very promising strategies in preventing cancer due to both additive and synergistic effects^[8]. Interest in the pharmacological effects of bioactive compounds on cancer treatments and prevention has increased dramatically over the past 20 y. Many fruits such as guava^[13], banana^[11], papaya^[12], lemon^[14], orange^[15], apple, water melon, litchi are reported to possess proven medicinal activities when

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms

*Address for correspondence
E-mail: psureshars@gmail.com

Accepted 30 March 2019
Revised 05 December 2018
Received 29 September 2018
Indian J Pharm Sci 2019;81(3):464-473

used as a whole fruit, seeds, leaves, and as peels with anticancer potential^[16-18].

Banana is one of the major fruit crops grown in India with the production of more than 30 million tons per annum. The *Nendran* banana is a dual-purpose banana, being utilized as a fruit and as in cooked form. Moreover, the chips, oil-fried products of *Nendran* increase the bioavailability of lipid soluble β -carotene in human diet, was reported by National Institute of Nutrition, India^[10]. The chips industry with the production of 2 lakh tons of chips and the value of 500 crores annually, generate equal volume of peel and the amount of waste is expected to increase with the development of clustered processing industries^[11]. Peel, the main by-product of banana constitutes 30-40 % of the fresh biomass of the *Nendran* fruit was found to contain much higher beneficial compounds compared to other fruit parts^[12,13]. They are rich source of crude protein (6-9 %), crude fat (3.8-11 %), total dietary fibre (43.2-49.7 %), polyunsaturated fatty acids, essential amino acids and micronutrients on dry weight basis. Banana peels are also a good source of lignin (6-12 %), pectin (10-21 %), cellulose (7.6-9.6 %), hemicelluloses (6.4-9.4 %) and galacturonic acid^[9]. The natural bioactive compounds such as carotenoids, quercetin derivatives, phenolic acids and saponins are found in the peels in high concentration^[14]. Like pulp, banana peel flour offers scope for making new processed nutraceutical products. These parts of banana have long been used in traditional medicine in Asia and Africa^[17]. Peel is used to make poultice for wounds to reduce pain and swelling. Peel, placed on a wart, with the yellow side out, could be an alternative to kill of wart and to reduce the irritation after a mosquito bite^[18,19]. Therefore, bioactive compounds from various parts of banana are considered to be natural free radical defence that were acknowledged to be beneficial for human health as an antioxidant, antitumor, and antimicrobial agent^[18,19].

Considering the need for effective anticancer agents, and the association of fruit and vegetable consumption with reduced cancer risk^[11,9,20-24], present study was conducted to profile the bioactive compounds in *Nendran* peel extract and to assess its antioxidant and anticancer activities with the focus to determine their ability to inhibit the proliferation of human breast cell line (MCF-7).

MATERIALS AND METHODS

The peel of the banana cv. *Nendran* was collected from the bakeries making banana chips in Tiruchirapalli,

Tamil Nadu and from the experimental farm of ICAR-National Research Centre for Banana, Trichy. The basic parameters such as fruit weight (g), length (cm), fruit diameter (cm), peel weight (g), pulp weight (g), firmness (kgf) and colour index were recorded. The peel and pulp were freeze-dried and ground into a fine powder using a mortar and pestle and passed using US 100 sieve (particle size 0.15 mm) and stored at 4° in a sealed poly propylene bag until further analysis.

Quantitative analysis of secondary metabolites:

Total carotenoids^[7], β -carotene^[8], total starch^[11], total and reducing sugar, protein^[9], superoxide dismutase (SOD)^[8], catalase (CAT), peroxidase (POX), glutathione reductase^[12], ascorbate peroxidase (APOX) and mineral nutrient^[8] were estimated using standard procedures. Fresh samples of peel and pulp were used for the estimation of phenols, flavonoids, secondary metabolites, minerals and other proximate analyses.

Estimation of total phenolic compounds:

Phenolic compounds from fruit peel and pulp extract were estimated according to the method outlined by Kumar *et al.*^[8]. In brief, 1 g of fresh sample was extracted using 80 % aqueous methanol and 1 ml of the sample was mixed with 1 ml of Folin-Ciocalteu phenol reagent and 1 ml of 20 % anhydrous sodium carbonate. The optical density was measured after 30 min at a wavelength of 650 nm. The phenolic content was obtained from a standard curve of gallic acid and then calculated as milligram phenolic per gram fresh weight of sample.

Estimation of total flavonoids:

Aluminium chloride colorimetric technique was used for flavonoid estimation^[10]. Fruit peel and pulp was extracted using 1 g/10 ml of 80 % aqueous methanol solvent. From the extract 0.5 ml was mixed with 1.5 ml of methanol, 0.1 ml of 10 % aluminium chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water, left at room temperature for 30 min and then the absorbance of the reaction mixture was measured at 416 nm with a double beam UV/Vis spectrophotometer (UV-3200, LabIndia). The total flavonoid content was obtained from a calibration curve, using quercetin as standard and then calculated as milligram flavonoids/g fresh weight of sample.

Diphenyl picrylhydrazyl (DPPH) radical scavenging assay:

DPPH stable free radical scavenging activity was

determined according to the protocol of Kumar *et al.*^[8] with a slight modification. Five grams of fresh sample was taken and extracted with 25 ml methanol. One millilitre of the extract was mixed with an equal volume of methanol to which 1 ml of 200 μ M DPPH was added. Control, blank and sample solutions were incubated in the dark at 37° for 30 min. Absorbance was measured at the end at 515 nm against control using a spectrophotometer. Ascorbic acid was used as the reference material and the percent inhibition was calculated by comparing the optical density values of the extract with those of the controls.

Ferric reducing antioxidant power (FRAP):

The antioxidant capacity of pulp and peel of the fruits were estimated spectrophotometrically following the procedure reported by Benzie and Strain^[25]. FRAP reagent was prepared by mixing 300 mM acetate buffer, 10 ml 2,4,6-tripyridyl-s-triazine in 40 mM HCl and 20 mM FeCl₃.6H₂O in the proportion of 10:1:1 at 37°. 3.995 ml of freshly prepared working FRAP reagent was pipetted and mixed with 5 μ l of the appropriately diluted plant sample and mixed thoroughly. An intense blue colour complex was formed and the absorbance at 593 nm was recorded against a reagent blank after 30 min incubation at 37°. The standard graph was plotted using FeSO₄.7H₂O at different concentrations. The reduction of Fe³⁺ into Fe²⁺ was observed by comparing the absorbance and the FRAP activity was expressed as μ M FeSO₄/ml. β -hydroxy anisole (BHA) was used as standard.

MTT assay for cell cytotoxicity:

Lyophilized peel powder (10 g) was extracted using 80 % methanol (1:3) and was concentrated to obtain 18 % aqueous methanol extract. The extract was used for further analysis of anticancer activity using MTT and acridine orange/ethidium bromide (AO/EB) assay. Human breast carcinoma (MCF-7) cell line was obtained from the National Centre for Cell Science, Mumbai, were cultured in liquid medium (Dulbecco's modified Eagle media, DMEM) supplemented with 10 % fetal bovine serum (FBS), 100 U/ml penicillin and 100 μ g/ml streptomycin and maintained under 5 % CO₂ atmosphere at 37°. Briefly, the cultured MCF-7 cells were harvested by trypsinization and pooled in a 15 ml tube^[12,26]. Then, the cells were plated at a density of 1 \times 10⁵ cells/well (200 μ l) into 96-well tissue culture plate in DMEM medium containing 10 % FBS and 1 % antibiotic solution for 24-48 h at 37°. The wells were washed with sterile PBS and treated with various

concentrations of 80 % aqueous methanol extract in a serum free DMEM medium. Then, the cells were incubated at 37° in a humidified 5 % CO₂ incubator for 24 h. After the incubation period, MTT (20 μ l of 5 mg/ml) was added into each well and the cells incubated for another 2-4 h until purple precipitate was clearly visible under an inverted microscope. Finally, the medium together with MTT (220 μ l) were aspirated off the wells and washed with 1X PBS (200 μ l). Furthermore, to dissolve formazan crystals, DMSO (100 μ l) was added and the plate was shaken for 5 min. The absorbance for each well was measured at 570 nm in a microplate reader (Thermo Fisher Scientific, USA), the percent cell viability and IC₅₀ values were calculated using GraphPad Prism 6.0 software (USA).

AO/EB staining:

AO/EB staining assay was performed as outlined by Ciniglia *et al.*^[24]. Briefly, 5 \times 10⁵ MCF-7 cells/ml were seeded into the 24 well tissue culture plate and treated with the best optimized extract in a serum-free DMEM medium. The plate was incubated at 37° in a 5 % CO₂ incubator for 24 h. After incubation, 50 μ l of 1 mg/ml AO and EB were added to the wells and mixed gently. Finally, the plate was centrifuged at 800 rpm for 2 min and evaluated immediately within 1 h and examined at least 100 cells by fluorescence microscope using a fluorescent filter. Early apoptosis, late apoptosis and necrosis indices were calculated using the difference in 100 cells with the staining^[24].

Statistical analyses:

Results were calculated as mean \pm standard deviation (SD) for each group and were analysed using GraphPad Prism (V.7, GraphPad Software, CA, USA) statistical software. For calculation of significant difference between multiple groups, t test was performed using SPSS (V.20.1, USA), one-way analysis of variance and post hoc analysis was performed with Tukey's test and p < 0.05 was considered statistically significant.

RESULTS AND DISCUSSION

The general physical parameters of the *Nendran* banana are presented in Table 1. *Nendran* a dual-purpose banana variety weighed between 140-225 g with the fruit length of 19.0-25.0 cm and the fruit circumference of 13 to 16.0 cm. The pulp weight varied between 97.5-124.67 g whereas the peel weighed in the range of 42.7-70.83 g. Colour, an important parameter for the selection of fruits expressed the L value of 43.6-52.4.

TABLE 1: PHYSICAL PARAMETERS OF NENDRAN BANANA (AAB)

Parameters	Values
Fruit weight (g)	140.8-225.34
Fruit length (cm)	18.5-24.3
Fruit circumference (cm)	12.5-15.4
Pulp weight (g)	97.5-124.67
Peel weight (g)	42.7-70.83
Firmness (kgf)	26.7-29.15
	L= 43.6-52.4
Colour index	a=16.21 b=21.9
Pulp to peel ratio	1.75-1.94

This has shown that the fruit is light in colour with the medium redness value. The peel to pulp ratio was 1.7-2.28, this explained the need for the study as huge biomass is generated as a waste while utilizing banana fruits in processing industry.

Chemical parameters of both peel and pulp were analysed and presented in Table 2. All the basic nutrient parameters like sugar, carbohydrate, protein and fat were more in the pulp than in the peel. The pulp contained higher value on total suspended solids (TSS) than the peel. Higher TSS was required for palatability of the fruit pulp and for higher consumer acceptance, while making processed products. Similarly, reducing sugars, non-reducing sugars and total carbohydrates were more in the pulp than in the peel. However, the peel gave higher value for fat and the ash content. The higher fat content in the peel facilitates its use for the extraction of essential oil. The higher ash content indicated higher mineral accumulation in the peel than in the pulp (Table 3). The nutrient composition of the cultivar is mostly depended upon its genetic makeup with a little influence of the environment and added nutrients^[11]. Accumulation of minerals is normally more in the peel than the pulp during fruit development stage. Among the minerals, potassium content was more with the values of 618.4 and 397.9 mg/100 g in the peel and the pulp, respectively followed by magnesium (536 and 421 ppm) and calcium (220 and 130 ppm). The positive effect of potassium on salt regulation and use of magnesium on reducing hypertension was well-documented^[3,17,21]. Consequently, these peels rich in minerals has a greater scope to be used in nutraceutical and cosmetic industries^[11,22,23] and as a supplement in bakery and biscuits units with its higher content of dietary fibre.

In the current investigation, aqueous methanol extract of *Nendran* peel showed highest flavonoid content of 51.47 µg/g and phenolic content of 0.54 mg/g, which

could be related to its antioxidant potential (Table 4). Medium containing polar and non-polar solvents has the capacity to release the cell wall bound polyphenols from the cells and neutralize the activity of poly phenol

TABLE 2: NUTRIENT COMPOSITION OF PULP AND PEEL OF NENDRAN BANANA (FWB)

Parameter	Peel	Pulp	p (0.05)	t-value (5 %)
Moisture (%)	75.3±3.2	62.0±2.87	1.24	2.34
TSS° Bx	6.4±0.25	21.5±0.81	0.43	-24.18
Reducing sugar (%)	12.6±0.47	17.8±0.64	0.57	-10.23
Non-reducing sugar (%)	3.4±0.19	5.4±0.09	0.67	-11.76
Starch (%)	4.5±0.11	7.6±0.19	0.28	-10.21
Protein (%)	0.35±0.004	0.83±0.04	0.23	-10.72
Fat (%)	3.67±0.20	2.58±0.12	0.64	09.27
Total carbohydrate (%)	22.6±1.62	32.5±1.99	0.42	-2.85
Ash (%)	1.83±0.10	1.54±0.07	0.26	1.61

(n=8)

TABLE 3: MINERAL CONTENT OF PULP AND PEEL SAMPLES OF NENDRAN (FWB)

Parameter	Peel	Pulp	p (0.05)	t-value (5%)
Potassium (mg/100 g)	618.4±25.74	397.9±15.05	0.43	13.11
Calcium (ppm)	220.0±08.32	130.0±04.68	0.57	13.76
Magnesium (ppm)	536.0±30.37	421.0±7.29	0.67	4.2
Iron (ppm)	12.4±0.32	4.7±0.11	0.29	17.59
Zinc (ppm)	2.8±0.06	1.4±0.06	0.56	11.43
Manganese (ppm)	2.3±0.07	1.3±0.05	0.63	18.8
Sodium (ppm)	38.0±2.91	22.0±1.34	0.42	3.95

(n=8)

TABLE 4: SECONDARY METABOLITES AND ANTIOXIDANT POTENTIAL OF PULP AND PEEL OF NENDRAN (FWB)

Parameter	Peel	Pulp	p (0.05)	t-value (5%)
Total phenols (mg/g)	0.54±0.02	0.82±0.03	0.42	-14.39
Total flavonoids (µg/g)	51.47±1.94	49.20±1.77	0.67	1.39
Total carotenoids (µg/g)	14.6± 0.33	15.8±0.78	0.56	-1.26
β carotene (µg/g)	10.3±0.33	12.6±0.59	0.63	-4.62
Ascorbate peroxidase (µmol/mg/min)	8.7±0.66	18.9±1.15	0.42	-5.91
Catalase (µmol/mg/min)	6.98±0.41	9.35±0.47	0.26	-2.77
Peroxidase (µmol/mg/min)	30.74±1.27	10.74±0.41	0.42	2.11
SOD (U/mg/min)	1.36±0.05	2.97±0.11	0.57	-18.92

(n=8)

oxidase, which generally degrades the polyphenols in plants^[21]. Phenols and flavonoids, the bioactive health beneficial phytochemicals, implicated in the prevention of a variety of diseases like cardiovascular diseases, cancers and neurodegenerative disorders was documented by researchers in different plant parts^[24-29].

FRAP assay (fig. 1a) is a widely used method that uses antioxidants as reductants in a redox-linked colorimetric reaction. The free radical scavenging activity was found better with peel than the pulp. With the concentration of 300 $\mu\text{g/ml}$, the peel has the antioxidant activity, matched with the BHA standard. Phenols and carotenoids were responsible for the higher FRAP activity in the peel. Carotenoids have good effect on human health, such as pro-vitamin A, antioxidant, anticancer, antiobesity effect and anabolic effect on bone components, which accumulated highly in pulp than peel^[20]. The higher carotenoid content of *Nendran*, make it as a nutraceutical variety and gives its intense yellow colour while ripening^[19].

The ability of the methanol extract of the *Nendran* peel to scavenge free radicals was determined based on its hydrogen donating ability in the DPPH assay (fig. 1b). The free radical scavenging activity of the extract was expressed as an IC_{50} of 180.33 $\mu\text{g/ml}$, which could be due to the presence of high flavonoid content, which might play a key role with its proton donating ability and could serve as free radical scavengers, acting possibly as primary antioxidants^[13]. Previous studies reported the antioxidant potential of extracts of banana pulp and peel with varying IC_{50} values^[13,29]. Similar to FRAP, in the present study, the DPPH free radical scavenging activity of the peel reached its peak at a concentration of 300 $\mu\text{g/ml}$ of extract and remained

constant at higher concentrations. The inhibition was almost linear in the case of pulp with highest activity observed at a dose of 500 $\mu\text{g/ml}$ of the extract. The variation in the antioxidant potential reported by several groups could be due to differences in cultivars, extraction procedures, geographical location and prevailing conditions such as soil, temperature, sunlight and horticultural practices^[4,7,9,11,30].

Similarly, the antioxidant enzyme concentration such as APOX, CAT, POX and SOD were found higher in the pulp than in the peel. These enzymes are the most important antioxidant enzymes scavenging the active oxygen species in plant cells^[16]. CAT may be major antioxidant enzyme involved in defence mechanism because it is induced during heating and persists during cold storage^[31]. POX, APOX and SOD participate in a great number of physiological processes, such as the biosynthesis of lignin and ethylene, defence against pathogens, wounding and stress response^[32]. POXs are recognized to be one of the most heat stable enzymes in plant and its resistance to heat has been reported by a numerous researchers^[7,9,11,30], which was more in peel than the pulp.

To know the effect of secondary metabolites on antioxidant assays, a correlation study was performed (fig. 2). Results indicated that with increasing concentration of the extract, the DPPH activity of both the peel and the pulp was increased. A linear correlation was observed with pulp on antioxidant activity whereas the peel exhibited the non-linear correlation with peak activity reached at a concentration of 300 $\mu\text{g/ml}$ of the extract. The FRAP activity also increased with increasing the concentration of the extract. Secondary metabolites exerted significant role in antioxidant




Fig. 1: Concentration-response of *Nendran* peel extract in FRAP and DPPH assays
 a. Concentration-response of *Nendran* extracts in FRAP assay ($\mu\text{M}/\text{FeSO}_4$) and (b) in DPPH assay (% inhibition) IC_{50} values of pulp (—■—) 432.72 $\mu\text{g/ml}$, peel (...●...) 180.33 $\mu\text{g/ml}$ and (...▲...) β -hydroxy anisole (BHA) as a standard




Fig. 2: Correlation between phenol and flavonoid content and antioxidant activity of *Nendran* banana
Correlation between total phenol and flavonoid content of pulp (—■—) and peel (...●...) of *Nendran* and antioxidant activity

activity. FRAP activity showed significantly positive correlation with total phenols and carotenoids whereas the DPPH activity was higher with the flavonoid content. This explained the fact that more antioxidant activity is required to study the full potential of extracts of natural products. Depending upon the results, further isolation and characterization could be decided.

The anticancer properties of the pulp with its tumour necrosis factor were already reported by various researchers^[11,19,23]. With higher content of phenols and flavonoids, the peel of *Nendran* bananas was chosen for evaluation of anticancer potential on MCF-7 cells. MCF-7 cell line has become a prominent model system for breast cancer studies as it relates to the susceptibility of the cells to apoptosis. Even though many tumours initially respond to chemotherapy, breast cancer cells can subsequently survive and develop resistance to the treatment. The inhibitory effect on breast cancer cells at different concentrations of peel extract of *Nendran* banana was studied. The dose-response relationship of the extract shown in Table 5. The intensity of dark-blue colour developed by the MTT reaction is directly proportional to the number of viable cancer cells. Cell

TABLE 5: EFFECT OF *NENDRAN* PEEL EXTRACT ON CELL VIABILITY AND CYTOTOXICITY OF MCF-7 CELL LINE

Extract concentration (µg/ml)	Cell viability (%)*	Cytotoxicity (%)
Control	100±0.00 (90.00)	0
200	24.7±1.10 (29.76)	75.3
180	48.08±0.86 (43.88)	51.92
160	55.02±1.24 (47.86)	44.98
140	56.78±0.64 (48.87)	43.22
120	59.73±1.48 (50.60)	40.27
100	67.58±0.58 (55.29)	32.42
80	70.72±0.45 (57.22)	29.28
60	74.26±0.97 (59.50)	25.74
40	80.52 ±0.96 (63.81)	19.48
20	91.14±2.34 (72.94)	8.86

*value in parenthesis are actual numbers recorded during the experiment (n=8)

viability (%) indicates the number of living cells. As the concentration of sample increased (µg/ml), the absorbance decreased that related to cell viability (%). As the concentration increased from 20 to 200 µg/ml, the cell viability of MCF-7 decreased from 91.14 to 24.7%. Cytotoxicity of the sample extract with varying concentration was also studied (Table 6). We noticed

TABLE 6: IC₅₀ VALUES AND REGRESSION COEFFICIENT OF CELL VIABILITY TEST WITH NENDRAN PEEL IN MTT ASSAY

Log(inhibitor) vs. normalized response -- Variable slope	
Best-fit values	
Log IC ₅₀	2.081
Hill Slope	-2.113
IC ₅₀	120.6
LogIC ₅₀	0.02057
Hill Slope	0.2733
95% Confidence intervals	
Log IC ₅₀	2.039 to 2.123
Hill Slope	-2.673 to -1.554
IC ₅₀	109.4 to 132.9
R square	0.85

that in control, mortality was not found on the cell population whereas with the concentration of 100 µg/ml, cytotoxicity of sample found to be 32.42 % while with 200 µg/ml of sample showed the mortality of 75.3 % of cancer cells, which implied the effectiveness of peel extract against the cancer. The half maximal inhibitory concentration (IC₅₀) is a measure of the effectiveness of a substance in inhibiting a specific biological or biochemical function. This quantitative measure indicates how much of inhibitor (sample) is required to inhibit a given biological process. In our present study, the extract of *Nendran* peel recorded IC₅₀ of 120.6 µg/ml. Therefore, it can be concluded that the effective concentration of the sample to arrest the cancer cell was 120.6 µg/ml, which recorded the regression coefficient value of 0.85. This explained the best fit of the model system followed for the experiment.

These changes were more visible in the dose response curve (fig. 3) plotted with OD value versus increasing concentration of the sample. The intensity of colour decreased with higher concentration of the extract with the values falling from 0.45 OD value at 570 nm in control to 0.05 OD value with the concentration of 200 µg/ml. The effect of peel extract on change in the cell viability was examined under the microscope (fig. 4) and the formazan crystals formed in the viable cell was noticed. The number of viable cells was more in untreated MCF-7 cells than in the *Nendran* peel-treated samples. This indicated the potential of peel as an anticancer agent and the presence of bio-active compounds, which had the capability to suppress the cell proliferation.

Tumor cells undergo apoptosis in the presence of anticancer drugs while normal cells become necrotic if the drug is toxic. MTT assays cannot differentiate

between these mechanisms of cell death. Therefore, the effects of drug may primarily be toxic or poisoning to normal cells. These drugs have limited clinical application^[11,31,32]. Therefore, detection of tumor cell apoptosis is more valuable than generally assessing tumour cell viability. To distinguish normal, early apoptotic, late apoptotic cells, and dead cells, nuclear morphology must be assessed. Fluorescent staining using AO alone has been used in the past; however, detection of cell apoptosis using AO/EB is a relatively new approach, and few papers have reported its use^[33-39]. Dual AO/EB fluorescent staining can detect basic morphological changes in apoptotic cells. In addition, it allows for the distinction between normal cells, early and late apoptotic cells, and necrotic cells. Therefore, AO/EB staining is a qualitative and quantitative method to detect apoptosis^[18].

Plant extract treated MCF-7 cells were subjected to AO/EB staining. AO will enter the nucleus and stain live cells as green colour fluorescence and EB will penetrate the nucleus of dead cells due to loss of membrane integrity and stain as red colour fluorescence (fig. 5a and 5b) when bound to concentrated DNA fragments or apoptotic bodies^[18,40]. In the present study, the MCF-7 cells were treated with 120.6 µg/ml of plant extract and the cells were examined using fluorescent microscopy. Normal viable cells appeared as green fluorescence with highly organized nuclei. Early apoptotic cells (28 %) were appeared as a crescent-shaped or granular yellow-green with AO nuclear staining. Late apoptotic cells (65 %) were appeared as a concentrated and asymmetrically localized orange nuclear ETBr staining^[33-36] (fig 6). Necrotic cells (7 %) showed uneven, orange-red fluorescence at their periphery without chromatin fragmentation. The IC₅₀ values of plant extracts-treated cells (120.6 µg/ml) showed typical apoptotic and necrotic morphological features such as condensed nuclei, membrane blebbing




Fig. 3: Dose response of *Nendran* peel extract on MCF-7 cells
Dose response curve of different concentrations (µg/ml) of *Nendran* peel extract on MCF-7 cells




Fig. 4: Effect of *Nendran* peel extract on the presence of viable cells through formation of formazan crystals after MTT treatment Influence of varying concentrations of *Nendran* peel extract on the presence of viable cells through formation of formazan crystals after MTT treatment (a) control, (b) cell behaviour at 40 µg/ml, (c) cell behaviour at 100 µg/ml and (d) cell behaviour at 200 µg/ml




Fig. 5: AO/EB staining of MCF-7 cell line treated with peel extract of *Nendran* banana Acridine orange/ethidium bromide (AO/EB) staining of MCF-7 cell lines (a) before treatment and (b) after treatment with peel extract of *Nendran* banana




Fig. 6: Effect of peel extract on early, late apoptosis and necrosis of MCF-7 cells (■) Early apoptosis, (□) late apoptosis, (▨) necrosis (%)

and formation of apoptotic bodies, which were clearly observed under the fluorescence microscope^[24,12]. These changes imply the cells appeared to be in the process of disintegrating. Whereas in control, the nucleus of the cells was regular in shape with no changes and retained the green colour. Our experiment result on anticancer activity of banana peel extract was corroborated with the earlier reports showing the similar efficacy with different parts like pulp^[11,39,40] and seed^[7,9,11] in inhibition of cancer cell lines.

The present investigation provides the gateway of information that aqueous methanol extract of *Nendran* peel has potent cytotoxic activity against MCF-7 breast

cell lines. Analysis of the extract showed the presence of high flavonoid content, a secondary metabolite, which is responsible for various therapeutic effects. The result obtained in this study clearly demonstrates that banana fruit peel has a broad spectrum of biological activities could be used as good source of antioxidant and antitumour agent. However, further research is needed to identify the bio-active components in banana peel extracts, their purification and characterization of compounds coupled with the evaluation of their mode of action to use the peel extracts in pharmaceutical industries.

Acknowledgments:

The authors thank the Director, ICAR-NRCB, Trichy for providing infrastructure facilities to carry out the research work. The authors also thank DBT for providing fund to carry out the work under North Eastern project. The Trichy research centre for Biotechnology Laboratory is also acknowledged for doing the antitumor assay. The technical help rendered by Dr. J. Jeyabaskaran and Pitchaimuthu is also acknowledged.

REFERENCES

- Mukherjee AK, Basu S, Sarkar N, Ghosh AC. Advances in cancer therapy with plant based natural products. *Curr Med Chem* 2001;8:1467-86.
- Lazarus D. Radiolarians decreased silicification as an evolutionary response to reduced Cenozoic ocean silica availability. *Proc Nat Acad Sci USA* 2009;106:9333-8.
- Mani RK, Kavitha K, Singh JD, Thakore P. A brief review of plants having anticancer property. *Int J Pharm Res* 2010;48:64-8.
- Shixin D, Brett J, West C, Jarakae J. A quantitative comparison of phytochemical components in global noni fruits and their commercial products. *Food Chem* 2010;31:935-8.
- Unnati S, Ripal S, Sanjeev A, Niyati A. Novel anticancer agents from plant sources. *Chin J Nat Med* 2013;11:16-23.
- Demain AL, Vaishnav P. Natural products for cancer chemotherapy. *Microb Biotechnol* 2011;4:687-99.
- Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr* 2004;134:3479-85.
- Kumar PS, Singh Y, Nangare DD, Bhagat K, Kumar M, Taware PB, *et al.* Influence of growth stage specific water stress on the yield, physico-chemical quality and functional characteristics of tomato grown in shallow basaltic soils. *Sci Hort* 2015;197C:261-71.
- Prasad KN, Hao J, Shi J, Liu T, Li J, Wei X. Antioxidant and anticancer activities of high pressure-assisted extract of longan (*Dimocarpus longan* Lour.) fruit pericarp. *Innov Food Sci Emerg* 2009;10:413-9.
- Praveena M, Prabha MS, Ravi I, Vaganan MM. Anti-colorectal cancer properties of Hill banana (cv. *Virupakshi* AAB) fruits: An *in vitro* assay. *Indian J Nat Sci* 2018;8(47):13226-33.
- Suresh Kumar P, Shiva KN, Mayil Vaganan M, Uma S. Waste utilization and functional foods from Banana. *Indian Hort* 2018;63(4):43-6.
- Zhang CR, Jayasree E, Suresh Kumar P, Nair MG. Antioxidant and Anti-inflammatory compounds in nutmeg (*Myristica fragrans*) pericarp as determined by *in vitro* assays. *Nat Prod Commun* 2015;10(8):1399-402.
- Rahmat A, Rosli R, Zain WN, Endrini S, Sani HA. Antiproliferative activity of pure lycopene compared to both extracted lycopenes and juices from watermelon (*Citrullus vulgaris*) and papaya (*Carica papaya*) on human breast and liver cancer cell lines. *J Med Sci* 2002;5:2097-103.
- Chen KC, Peng CC, Chiu WT, Cheng YT, Huang GT, Hsieh CL. Action mechanism and signal pathways of *Psidium guajava* L. aqueous extract in killing prostate cancer LNCaP cells. *Nutr Cancer* 2010;62:260-70.
- Miller EG, Porter JL, Binnie WH, Guo IY, Hasegawa S. Further studies on the anticancer activity of citrus limonoids. *J Agric Food Chem* 2004;60:69-74.
- Arora M, Kaur P. Antimicrobial & antioxidant activity of orange pulp and peel. *Int J Sci Res* 2013;2:412-5.
- Manosroi J, Dhumtanom P, Manosroi A. Anti-proliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *Cancer Lett* 2006;235:114-20.
- Tsamo CVP, Herent M, Tomekpe K, Emaga TH, Quetin-Leclercq J, Rogez H, *et al.* Phenolic profiling in the pulp and peel of nine plantain cultivars (*Musa* sp.). *Food Chem* 2015;15:197-204.
- Orhue PO, Momoh AR. Antibacterial activities of different solvent extracts of *Carica papaya* fruit parts on some gram positive and gram-negative organisms. *Int J Herbs Pharmacol Res* 2013;2:42-7.
- Kumar KPS, Bhowmik D, Duraivel S, Umadevi M. Traditional and medicinal uses of banana. *J Pharmacogn Phytochem* 2012;1:51-63.
- Emaga TH, Andrianaivo RH, Wathelet B, Paquot M. Effects of the stage of maturation and varieties on the chemical composition of banana and plantain peels. *Food Chem* 2007;103:590-600.
- Ighodaro O. Evaluation study on Nigerian species of *Musa paradisiaca* peels: phytochemical screening, proximate analysis, mineral composition and antimicrobial activities. *Researcher* 2012;4:17-20.
- Li ZY, Wang Y, Shen WT, Zhou P. Content determination of benzyl glucosinolate and anti-cancer activity of its hydrolysis product in *Carica papaya* L. *Asian Pac J Trop Med* 2012;72:382-5.
- Ciniglia C, Pinto G, Sansone C, Pollio A. Acridine orange/Ethidium bromide double staining test: A simple *In-vitro* assay to detect apoptosis induced by phenolic compounds in plant cells. *Allelopathy J* 2010;26(2):301-8.
- Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay. *Anal Biochem* 1996;239:70-6.
- Houston MC. Treatment of Hypertension with Nutraceuticals, Vitamins, Antioxidants and Minerals. *Expert Rev Cardiovasc Ther* 2007;5:681-91.
- Karppanen H. Minerals and Blood pressure. *Ann Med* 1991;23:299-305.
- Donovan JL, Meyer AS, Waterhouse AL. Phenolic composition and antioxidant activity of prunes and prune juice (*Prunus domestica*). *J Agric Food Chem* 1998;46:1247-52.

29. Ediriweera ER, Rathnasooriya WD. A review of herbs used in diabetes mellitus by Sri Lankan Ayurvedic and traditional physicians. *Ayu* 2009;30:373-91.
 30. Freitas D, Glories Y. Concentration and compositional changes of procyanidins in grape seeds and skin of white *Vitis vinifera* varieties. *J Sci Food Agric* 1999;79:1601-6.
 31. Ghasemi K, Ghasemi Y, Ebrahimzadeh MA. Antioxidant activity, phenol and flavonoid contents of 13 citrus species peels and tissues. *Pak J Pharm Sci* 2009;22:277-81.
 32. González-Gallego J, García-Mediavilla MV, Sánchez-Campos S, Tuñón MJ. Fruit polyphenols, immunity and inflammation. *Br J Nutr* 2010;104:15-27.
 33. Goulas V, Manganaris G. Exploring the phytochemical content and the antioxidant potential of Citrus fruits grown in Cyprus. *Food Chem* 2012;131:39-47.
 34. Huang ME. Oxygen metabolism and reactive oxygen species cause chromosomal rearrangements and cell death. *Microb Biotechnol* 2009;104:9747-52.
 35. Kim H, Moon JY, Kim H, Lee DS, Cho M, Choi HK. Antioxidant and antiproliferative activities of mango (*Mangifera indica* L.) flesh and peel. *Food Chem* 2010;121:429-36.
 36. Maria GA, Sonia PT, Celestino SB, Julian CRG. Evaluation of the antioxidant properties of fruits. *Food Chem* 2004;50:889-92.
 37. Lecoeur H. Nuclear apoptosis detection by flow cytometry: influence of endogenous endonucleases. *Exp Cell Res* 2002;277:1-14.
 38. Burda S, Oleszek W, Lee CY. Phenolic compounds and their changes in apples during maturation and cold storage. *J Agric Food Chem* 1990;38:945-8.
 39. Senthilraja P, Kathiresan K. *In vitro* cytotoxicity MTT assay in Vero, HepG2 and MCF -7 cell lines study of Marine Yeast. *J Appl Pharm Sci* 2015;55:689-94.
 40. Sumathy N, Sumathy J. Antibacterial and antifungal activity of musa fruit peels against skin and gastrointestinal tract diseases. *Herbal Tech Ind* 2011;4:11977-83.
-