

A Review on Chemical Composition, Oil Quality, and Bioactivity of Vetiver Essential Oil

A. PANDEY AND S.C. TIWARI*

Department of Forestry, Wildlife and Environmental Sciences, Guru Ghasidas Vishwavidyalaya, Bilaspur, Chhattisgarh 495009, India

Pandey *et al.*: Composition, Quality and Bioactivity of Vetiver Oil

Chrysopogon zizanioides is the well-known eco-friendly plant known for the essential oil obtained from its roots. It is the most popular essential oil which does not have any synthetic substitute available. The essential oil of vetiver is widely utilized in traditional Thai, Chinese, and Ayurvedic medicine to cure a variety of illnesses. The essential oils with various biological properties are employed in the culinary, pharmaceutical, food and beverage and cosmetic industries. The current study is the review of chemical makeup, oil quality (as per Bureau of Indian Standards) as well as the bioactivities the of the vetiver essential oil. Various pharmacological studies have found that vetiver oil has significant effects as an antifungal, antibacterial, antioxidant, anti-inflammatory, and cytotoxic agent against cancer, as well as being beneficial in aromatherapy. The essential oil used in conventional medicine to cure a variety of illnesses, such as mouth ulcers, fever, headaches, depression, nervous tension, anxiety disorders, insomnia, and sexual dysfunction. Root is used as alexiteric, stomachic, carminative, astringent, and boosts the immune system in addition to treating skin conditions, urinary problems, jaundice, and flatulence as well as anaemia, and haemoptysis.

Key words: Aromatic plant, essential oil, herbal medicine, vetiver

Essential Oils (EOs) do not need an introduction because they are highly regarded as plant-based products with enormous therapeutic and economic potential. They are obtained from various plant parts such as flowers, buds, stems, leaves, fruits, seeds, and roots, to name a few^[1]. Medicinal and aromatic plants are among the most important types of horticulture crops that play a special role in sustaining various pharmaceutical, perfumery, and cosmetic industries in India and abroad^[2-4]. The root is the most economically important part of the vetiver plant^[5].

Vetiver, botanically known as *Chrysopogon zizanioides* (*C. zizanioides*) (L.) Roberty Syn., *Vetiveria zizanioides* (*V. zizanioides*) Nash is a perennial herbaceous grass belonging to the Poaceae family. It is commonly known as Khus or Khas-Khas^[6,7]. Its origin is India, and it grows wildly in most parts of the world, from tropical to Mediterranean climates, including regions in Asia, Central and South America, Oceania, and Africa^[8-10]. It is grown for its exceptional ability to create an EO, a complex blend of sesquiterpene alcohols and hydrocarbons^[11,12]. Due to its distinctive aroma, which is not produced by any

synthetic chemical, vetiver oil is also known as the "oil of tranquility"^[11-14]. The largest producers of vetiver EO in the international market are Indonesia, Haiti, Brazil, China, India and Guatemala^[15-17,10]. The major consumers of vetiver oil are the United States, Europe, India, and Japan^[1,18]. In India, vetiver is mostly grown in the southern and peninsular regions, including Uttar Pradesh, Rajasthan, Bihar, Kerala, Tamil Nadu, Karnataka, and other states. There are two strains, South Indian and North Indian, with several vernacular names^[19,20,1,4]. India produces only 20-25 tonnes of vetiver EO annually, compared to the estimated 600-700 tonnes produced worldwide. The estimated cultivation area for vetiver has reached nearly 10 000 hectares^[10,21]. Vetiver grass is grown for the extraction of an EO with significant commercial value, used in fragrance and aromatherapy^[22,23]. It is mostly employed in high-end perfumery, where

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms

*Address for correspondence
E-mail: ankitforestry21@gmail.com

Accepted 26 August 2024
Revised 17 May 2024
Received 04 October 2023
Indian J Pharm Sci 2024;86(4):1176-1186

its odor persistence makes it valuable as a fixative when combined with other scents^[24,25]. The highly complex chemical makeup of vetiver oil consists primarily of sesquiterpenes and its derivatives, of which vetiverols, their carbonyl compounds, and esters are the primary components. The relative abundance of these chemicals determines the oil's quality. The main odour-influencing components in vetiver oils from various sources are three carbonyl compounds, Alpha (α)-vetivone, Beta (β)-vetivone, and khusimone^[2,26]. β -vetivone has a better odor and is regarded as the most important component, while its major isomer, nordihydro β -vetivone, has a strong, rich, woody-peppery note^[19,25]. These numerous active components of vetiver oil provide biological properties like antibacterial, antifungal, antiviral, antioxidant, anti-inflammatory, and anticancer properties^[25,13,15]. This review focuses on describing the changes in and availability of the chemical composition of vetiver oil from different geographic regions, the quality of the oil, and its uses in various pharmaceutical (bioactivity) and medicinal capabilities of vetiver EOs.

METHODS

An extensive search of scientific databases including the Web of Science, Google Scholar, Scopus, ISC, and PubMed was used in this review to gather evidence on vetiver's chemical

composition and phytochemical qualities. The keywords used in the database for the literature survey were chemical composition of vetiver, vetiver oil quality, pharmacological uses of vetiver, chemistry of vetiver oil, and bioactivity. The literature in this study has included original research papers, review papers, as well as a few books on herbal and traditional medicines.

EO EXTRACTION:

EOs produced from different herbs and aromatic plants have seen an increase in popularity in recent years. Hydro Distillation (HD) (traditional method), steam distillation, solvent extraction, microwave assisted extraction, Supercritical Fluid Extraction (SFE) and CO₂ expanded ethanol method are the current vetiver oil extraction techniques^[27,28,17,14]. In order to obtain the EOs from medicinal herbs and plants, HD has been the most popular technique among the various methods of oil extraction^[17]. The EO obtained from various extraction methods shows significant differences in yield percentage, chemical composition, extraction time, and other biological activities of vetiver oil, as presented in Table 1. Pripdeevech *et al.*^[29] reported that there was no discernible difference between the composition of vetiver oil obtained using the steam distillation and solvent extraction apparatus.

TABLE 1: GENERAL COMPARISON OF VETIVER OIL OBTAINED FROM DIFFERENT EXTRACTION METHODS

Parameter	Methods of oil extraction		
	HD	CXE	SFE
Yield (%)	0.6	5.12	0.5
Oil colour	Pale yellow	Light yellow	Greenish brown
Time of extraction (h)	12	3	2
Chemical composition			
Hydrocarbon	90.71	64.46	81.91
Alcohol	2.77	-	-
Carbonyl compound	-	5.42	-
Carboxylic acid	2.66	-	-
Total identified (%)	96.14	69.88	81.91

Note: CXE: Carbon dioxide expanded ethanol extraction

CHEMICAL COMPOSITION OF VETIVER EO:

Numerous studies have been conducted on vetiver oil chemical makeup. Over 300 chemicals have been identified, with sesquiterpenes and their derivatives like alcohols, hydrocarbons, and ketones dominating the majority of them^[30]. The oil produced by vetiver grown in regular soil had 37 volatiles, according to analysis. Pripdeevech *et al.*^[29] studied vetiver cultivation using different methods and found that 39 volatile components were present in the oil extracted from vetiver grown in normal soil with microorganisms. The major component was (Z)-9,10-dehydro-2-norzizaene, comprising 46.03 % of the oil. The oil produced by vetiver grown in normal soil had 37 volatiles, the three main ones among these were (Z)-9,10-dehydro-2-norzizaene (20.78 %), khusimone (20.57 %), and khusimol (11.11 %). Champagnat *et al.*^[30] reported that the EOs of *V. zizanioides* Nash from 9 nations-Brazil, China, Haiti, India, Java, Madagascar, Mexico, Reunion, and Salvador have different chemical compositions. About 110 components, mostly sesquiterpenes, were found in oils after analyses using Gas Chromatography-Mass Spectrometry (GC-MS). After the GC-MS analysis of vetiver oil, Hammam *et al.*^[31] found that the 15 different compounds shows that sesquiterpenes are the most common class of chemicals. The most crucial ingredients were alcohols and acids. The

main compounds among the alcohols in vetiver oil include preziza-7(15)-en-12-ol, cedren-5-en-15-ol, preziza-7(15)-en-3-ol, ziza-6(13)-en-12-ol (khusiranol), ziza-5-en-12-ol, 12-nor-ziza-6(13)-en-2-ol, and khusian-2-ol (helifolan-ol). Similarly, Oliveira *et al.*^[13] and Lunz *et al.*^[7] reported that there were 22 different compounds were found, the majority of which (70.4 %±0.1 %) are oxygenated sesquiterpenes. The primary ingredients in vetiver EO were khusimol (30.0 %±0.3 %), eudesmol (10.8 %±0.3 %), muurolene (6.0 %±0.1 %), and patchouli alcohol (5.6 %±0.2 %). The Srilanka ecotype of vetiver from Northeast Thailand was steam-distilled to produce a viscous light-brown oil with a balsamic, earthy, and sweet woody scent that contained between 0.3 and 1.0 % volume by weight. The main components were khusimol (12.7 %), longipinene (4.2 %), valeranol (3.9 %), and epizizanal (3.3 %), with α -vetivone and β -vetivone following closely behind (2.0 % and 1.6 %, respectively)^[32]. Similarly Lima *et al.*^[33] found that after the GC-MS analysis revealed a variety of substances, including khusimol (19.57 %), (E)-isovalencenol (13.24 %), α -vetivone (5.25 %), vetiselinenol (5.08 %), α -cadinol (5.01 %), α -vetivone (4.87 %), and hydroxy-valencene (4.64 %) being the main substances in the EO. The chemical composition of *C. zizanioides* EO with number of compounds has been found, compiled from the different geographic location of the world is shown in Table 2.

TABLE 2: CHEMICAL COMPOSITION OF VETIVER GRASS FROM THE DIFFERENT LOCATION OF WORLD STUDIED BY VARIOUS RESEARCHERS

S. No	Location	Chemical composition	Total no. of compounds	References
1	China	2,3,5,5,8,8-hexamethyl-cycloocta-1,3,6-triene, 1,5,9,9-Tetramethyl-2-methylene-spiro[3.5]non-5-ene, (+)-sativene, 4,8,8-trimethyl-2-methylene-4-vinylbicyclo [5.2.0]nonane, α -amorphene, 2-isopropenyl-1,3,5-trimethylbenzene, α -gurjunene, β -vatirenene, 9 d-cadinene, β -guaiane, dehydro-aromadendrene, cubenol, (+)-ledene, epiglobulol, widdrol, 6-Isopropenyl-4,8a-dimethyl-1,2,3,5,6,7,8,8a-octahydro-naphthalen-2-ol, 3-(2-isopropyl-5-methylphenyl)-2-methylpropionic acid, cedr-8-en-13-ol, ethyl 4-(4-methylphenyl)-4-pentenoate, isovellerdiol, α -curcumene, 3,3,8,8-tetramethyl-tricyclo[5.1.0.0(2,4)] oct-5-ene-5-propanoic acid, solavetivone, 3,8-dimethyl-4-(1-methylethylidene)-2,4,6,7,8,8a-hexahydro-5(1H)-azulenone, (-)-spatulanol.	25	[11]
2	Turkey	8,9-dehydro-cycloisolongifolene, cadina-1(10),4-diene, α -curcumene, γ -cadinene, β -vetivenene, γ -gurjunene, ledene oxide, η -himachalene, valencene, C ₁₅ H ₂₂ O, γ -muurolene, β -guaiane, allaromadendrene oxide, δ -cadinol, junipene, tau-murolol, trans-caryophyllene, cubenol, caryophyllene oxide, khusinol, 8-cedren-13-ol, C ₁₃ H ₁₈ O, β -gurjunene, thujopsene-13, dehydro-aromadendrene, β -vetivone, α -vetivone.	30	[26]

3	Comoros	<p>α-phellandrene, (E)-isoeugenol, khusimene, ar-curcumene, α-amorphene, valencene, γ-vetivenene, elemol, β-vetivenene, β-calacorene, spathulenol, trans-sesquisabinene hydrate, khusimone, 1,10-epi-cubenol, 10-epi-γ-eudesmol, 1-epi-cubenol, epi-α-cadinol, α-cadinol, 7-epi-α-eudesmol, valerianol, epizananone, khusinol, khusian-2-ol, zizanal, (Z)-β-santalol, nootkatol, juniper camphor, vetiselinenol, khusimol, (E)-β-santalol, 14-hydroxy-γ-cadinene, 13-hydroxy valencene, bicyclo- vetivenol, isovalencenol, nootaktone, β-vetivone, khusinol acetate, (Z)-β-santalol acetate, α-vetivone, epi-laurenene.</p>	41	[34]
4	Brazil	<p>Naphthalene, β-cubebene, Isolongifolene, β-gurjunene, α-patchoulene, β-cadinene, α-muurolene, α-amorphene, valencene, α-bulnesene, α-elemol, β-vatirenene, β-eudesmol, spathulenol, guaiol, humulane-1,6-dien-3-ol, cubenol, agarospirol, cedren-13-ol, patchouli alcohol, khusimol, nootkatone,</p>	22	[13]
5	India	<p>Cycloisolongifolene, all-oaromadendrene, α-copaene, γ-muurolene, germacrene D, guaia-6,9-diene, isolongifolol, isospathulenol, isoledene, epiglobulol, tau-muurolol, cubenol, isoaromadendrene epoxide, globulol, humulane-1,6-dien-3-ol, caryophyllene, aristolene, α-Ylangene, 2(1H) naphthalenone, 3,5,6,7,8,8a-hexahydro-4,8a dimethyl-6-(1-methylethenyl)-, β-vatirenene, α-cedrene, selin-6-en-4α-ol, γ-himachalene, valerenol, α-Guaiene, α-gurjunene, longiverbenone, α-longipinene, α-amorphene, β-copaene, nootkatone, isolongifolene, 4,5-dehydro-</p>	32	[18]
6	Thailand	<p>Tricyclene, camphene, camphor, bornyl acetate, 7-epi-Silphiperfol-5-ene, presilphiperfol-7-ene, silphiperfol-5-ene, silphiperfol-4,7(14)-diene, silphiperfol-6-ene, cadina-1(10), 6,8-triene, (+)-β-cedrene, caryophyllene(E-), β-copaene, 9-methyl-5-octahydroanthracene, α-humulene, cis-cadina-1(6),4-diene, α-neocallitropsene. Cuparene, 3-methyl-2-pent-2-enyl-cyclopent-2-enone, Cameroon-7-α-ol, Humulane-1,6-dien-3-ol, Silphiperfolan-6-β-ol, guaia-3,9-diene, prenopsan-8-ol, caryophyllene oxide, alloaromadendrene oxide-(2), humulene epoxide II, β-humulene, silphiperfol-6-en-5-one, α-acorenol, β-vatirenene, epi-β-bisabolol, 11αH-Himachal-4-en-1β-ol, 1,4-methanoazulen-7(1H)-one, octahydro-4,8,8,9-tetramethyl</p>	34	[35]
7	Taiwan	<p>2,3,5,5,8,8-hexamethyl-cycloocta-1,3,6-triene, 1,5,9,9-tetramethyl-2-methylene-spiro[3.5]non-5-ene, (+)-sativin, 4,8,8-trimethyl-2-methylene-4-vinylbicyclo[5.2.0]nonane, α-amorphene, 2-isopropenyl-1,3,5-trimethylbenzene, α-gurjunene, β-vatirenene, δ-cadinene, β-guaiene, dehydroaromadendrene, cubenol, (+)-ledene. epiglobulol, widdrol, 6-isopropenyl-4,8a-dimethyl-1,2,3,5,6,7,8,8a-octahydro-naphthalen-2-ol, 3-(2-isopropyl-5-methylphenyl)-2-methylpropionic acid, cedr-8-en-13-ol, ethyl 4-(4-methylphenyl)-4-pentenoate, isovellerdiol, α-curcumene, 3,3,8,8-tetramethyl-tricyclo [5.1.0.0(2,4)] oct-5-ene-5-propanoic acid, solavetivone, 3,8-dimethyl-4-(1-methylethylidene)-2,4,6,7,8,8a-hexahydro-5(1H)-azulenone, (-)-spathulenol</p>	25	[23]
8	Egypt	<p>Linalool, trans-rose oxide, menthone, trans methone, citronellol, D-Carvone, geraniol, citronellyl formate, geranyl formate, zizanal, β-bourbonene, acora-3(7),14-diene, caryophyllene, daucene, (+)-epi-bicyclo-sesquiphellandrene, prezizaene, khusimene, α-gurjunene, (1R,5R)-1-iso propyl-8-methyl-4-methyl lenespiro [4.5] dec-7-ene, selina-3,7(11)-diene, α-amorphene, cis-eudesma-6,11-diene, β-vetispirene, β-cadinene, γ-muurolene, D-cadinene, iso ledene, cadina-1(10),4-diene D-amorphene, 11,12,13-tris-nor-trans-eud esm-5-en-, 7-one, α-calacorene, β-vatirenene, isolongifolene, 4,5,9,10-dehydro-, 4-(1,3,3-trimethyl-bicyclo.[4.1.0]hept-2-yl)-but-3-en-2-one, γ-vetivenene, ylangene, β-maaliene, 13-nor-eremophil-1(10)-en-11-one, junenol, γ-selinene, β-cadinene, valencene, β-guaiene, 2-isopropyl-5-methyl-9-methylene-bicyclo-1-decene (4.4.0), cyclocopacamphenol, spiro [4.5] dec-8-en-7-ol, 4, 8-dimethyl-1-(1-methyl ethyl)-, zizanol, khusiol, juniper camphor, cycloheptane, 4-methylene-1-methyl-2-(2-methyl-1-propen-1-yl)-1-vinyl-, δ-selinene, vetiselinenol, α-isonootkatol, khusimol, α-vetivol, α-costol, valerenol, β-vetivenene, (e)-eremophila-1(10),7(11)-dien-12-ol (isoval encenol), (z)-isovalencenal, β-vetivone, (e)-isovalencenal eremophila-1(10),7 (11)-dien-12-al, α-vetivone</p>	63	[36]

9	Indonesia	<p>Khusimol (zizanol), vetiselinenol (isonootkatol), cyclocopacamphan-12-ol (epimer a), α-cadinol, α-vetivone (isonootkatone), β-vetivenene, β-eudesmol, β-vetivone, khusenic acid, β-vetispiroene, γ-vetivenene, α-amorphene, (e)-eudesm-4(15),7- dien-12-ol, b-calacorene, g-cadinene, (z)-eudesm-6-en-11-ol, g-amorphene, ziza-5-en-12-ol, b-selinene, (z)-eudesma-6,11-diene, salvia-4(14)-en-1-one, khusinol</p> <p>Cyclocopacamphan-12-ol (epimer b), selina-6-en-4-ol, khusian-ol, d-amorphene, 1-epi-cubenol, khusimene (ziza-6(13)- ene), ziza-6(13)-en-3b-ol, ziza-6(13)-en-3-one, 2-epi-ziza-6(13)-en-3a-ol, 12-nor-ziza-6(13)-en-2bol, a-vetispiroene, eremophila-1(10),7(11)- diene, dimethyl-6,7-bicyclo- [4.4.0]-deca-10-en-one, 10-epi-g-eudesmol, a-calacorene, (e)-opposita4(15),7(11)-dien12-ol, prekhuseniacid, 13-nor-eudesma-4,6-dien-11-one, isovalencenol, spirovetiva-1(10),7(11)-diene, 2-epi-ziza-6(13)-en-12-al, (e)-isovalencenal, preziza-7(15)-ene, (z)-eudesma-6,11-dien3b-ol, intermedeol (eudesm-11- en-4-ol), isoeugenol, isokhusenic acid, elemol, eremophila-1(10),6-dien12-al, juniper camphor, khusimone, eremophila-1(10),4(15)-dien-2a-ol, eremophila-1(10),7(11)-dien-2b-ol, (z)-isovalencenal, allo-khusiol, methyl-(e)-eremophila1(10),7(11)-dien-12-ether, (e)-2-nor-zizaene, (z)-eudesm-6-en-12-al, funebran-15-al</p>	61	[37]
---	-----------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----	------

An EOs aroma and chemical makeup can differ depending on the region, climate, and growing conditions (such as soil type, altitude, and water availability), the season (such as before or after the flowering period), and even the time of day when it is harvested^[1,25,15]. After studying the vetiver oil composition from various countries, as shown in Table 2, it was found that the highest number of compounds were present in Egypt, followed by Indonesia, while the lowest number of compounds were found in Turkey, with only 22 compounds. India has two recognised vetiver varieties: The Bharatpur or North Indian type is primarily found in the Indian states of Rajasthan, Madhya Pradesh, Uttar Pradesh, and Bihar. It has abundant flowering, high seed setting, narrow leaves, and vigorous roots that produce laevorotatory root oil (khush oil) with a low concentration and superior quality. The South Indian cultivated variety can be found in the states of Andhra Pradesh, Karnataka, Tamil Nadu, and Kerala along the East and West coastlines of the Indian peninsula^[1,7]. Chahal *et al.*^[1] conducted a study on Indian vetiver oil and reported that a total of 29 and 33 compounds in South and North Indian vetiver oils, respectively. The chemical composition of North Indian and South Indian strains of vetiver presented in Table 3.

The antipodal sesquiterpenes of the cadinane class are abundant in North Indian vetiver, while the α and β -vetivones are lacking. The dextrorotation of the south Indian vetiver root oil is visible. It contains the same sesquiterpene ketones, such as vetivones, vetispiroenes, eudesmane, and zizaane

sesquiterpenes that are present in vetiver root oils produced in other countries^[16]. Mallavarapu *et al.*^[16] found that 80 components have been identified, accounting for 94.5 %-97.8 % of the oils. Sesquiterpenes, oxygenated sesquiterpenes, and oxygenated sesquiterpenes with cedrane, bisabolane, eudesmane, eremophilane, and zizaane skeletons were abundant in the oils from Bangalore, Hyderabad, Kundapur, and Mettupalayam^[34-38].

PHYSICO-CHEMICAL PROPERTIES OF VETIVER EO:

Physico-chemical characteristics of the oil, including an assessment of its aroma, were used to determine its quality. The presence of several ketonic sesquiterpenes, including vetiverol, α -vetivone, and β -vetivone, as well as the ester vetiveryl vetivenate, gives vetiver oil its distinctive aroma. The fixative activity of oil is aided by sesquiterpene alcohols such as vetivol acetate and vetiverol. In a study by Kadarohman *et al.*^[39] vetiver EO quality was observed from organic and non-organic vetiver plants and found that compared to oil obtained from inorganic cultivation, organic was higher in quality. In terms of appearance, organic vetiver oil was more concentrated and had no black spots. Chemically, vetiver EO from organic cultivation includes more vetiverol and less pesticide residue^[40]. The quality of vetiver oil from different geographic locations as compared to Bureau of Indian Standards (BIS) and Karnataka Soaps and Detergents Limited (KSDL) value has been presented in Table 4.

TABLE 3: CHEMICAL COMPOSITION OF NORTH AND SOUTH INDIAN STRAINS OF VETIVER

S. No	Component	Concentration percent	
		North India	South India
1	β -caryophyllene	-	0.37
2	Sativen	0.65	-
3	Khusimene	0.66	1.59
4	α -humulene	-	0.03
5	α -amorphene	0.22	1.68
6	α -bisabolol	0.19	-
7	γ -cadinene	0.82	-
8	β -caryophyllene oxide	0.28	-
9	Allo-aromadendrene	0.17	-
10	Cubenol	1.64	-
11	δ -guaiene	-	1.42
12	Valencene	-	1.31
13	β -vetivenene	-	0.14
14	β -vatirenene	-	0.72
15	Caryophyllenyl alcohol	-	2.3
16	Humulane-1,6-dien-3-ol	0.87	1.42
17	10-epi- γ -eudesmol	0.46	-
18	β -lonol	0.46	-
19	β -bisabolol	0.57	1
20	Epiglobulol	1.95	-
21	Selina-6-en-4ol	-	2.11
22	Globulol	-	1.91
23	Junipene	5.54	1.6
24	α -cadinol	1.46	1.74
25	Guaiol	1.03	1.58
26	α -eudesmol	0.49	-
27	β -eudesmol	0.97	1.28
28	Allo-aromadendren oxide	-	0.83
29	β -cedren-9- α -ol	1.76	-
30	Farnesol	2.65	-
31	Khusinol	10.28	-
32	γ -muurolene	4.56	-
33	E-caryophyllene	0.74	-
34	γ -cadinene aldehyde	1.28	-
35	Patchouli alcohol	-	1.61
36	γ -gurjunepoxide	-	1.14
37	Cyclosativene	-	2.06
38	Cedryl acetate	-	0.29
39	Spathulene	1.28	2.37
40	γ -maaline	1.57	-
41	Calerene	1.57	-
41	γ -himachalene	0.23	-
42	Viridiflorene	-	4.64
43	Khusimol	16.25	15.77
44	Bicyclovetivenol	2.88	10.76
45	α -vetivone	2.01	3.04
46	Velerenol	-	-3.44
47	Germacrene-D	9.73	-
48	Cedrene-13-ol-8	2.23	1.05
	Total	33	29

TABLE 4: PHYSICO-CHEMICAL PROPERTIES OF VETIVER OIL

S.No	Parameter	BIS value	KSDL value	Putrawan et al. ^[40]	Kadarohman et al. ^[39]	Dubey et al. ^[38]	Khalil et al. ^[41]	Yanto et al. ^[2]
1	Colour	Brown to reddish viscous liquid	Brown to reddish viscous liquid	Pale yellow to reddish brown	Yellow to reddish brown	Brown to yellowish brown clear liquid	-	Reddish brown
2	Specific gravity	0.9926-1.0444	0.985-1.020	0.980-1.003	0.980-1.003	1.0118-1.0026	1.002	1.0097
3	Refractive index@ 24°	1.588-1.5306	1.5132-1.5242	1.52-1.53 @20°	1.520-1.530	1.5147-1.5220	1.5112	1.5308
4	Optical rotation		+15° to +35°		-	-57.70 to +28.23	+40.0	-
5	Acid value	8.4-40.1	35	10-35	-	21.30-28.45	46	11.5
6	Ester value	5.6-24.6	5-16	--	5-26	8.36-11.27	14	18.57
7	Saponification value	-	-	-	-	--	60	-
8	Odour	Rosy odour	Woody aroma	Vetiver characteristics		Heavy earthy, green to heavy woody	-	Slightly smoky

BIOACTIVITY OF VETIVER EO:

Antimicrobial activity:

The vetiver EO had excellent antibacterial efficacy against all of the clinical pathogens that were tested, with no discernible difference in susceptibility between gram-positive and gram-negative bacteria^[41-43]. By measuring the zone of inhibition, ethanolic root extract of *V. zizanioides* is shown to have antibacterial activity. The ethanolic root extracts exhibit antimicrobial properties due to presence of flavonoids, terpenoids, and tannins included in vetiver^[44]. Sangeetha et al.^[45] obtained similar findings and found that vetiver leaf extract had the highest mean zone of inhibition against *Staphylococcus aureus*. Soni et al.^[46] studied that vetiver solvent extracts have the potential to be utilised as an antimicrobial medication against infectious agents and as a treatment for a variety of infectious disorders. According to the Efe et al.^[43] findings, vetiver EO showed an inhibitory impact against the tested microorganisms that was greater than or comparable to that of antibiotics at high vetiver EO concentrations. In a study conducted by Nantachit et al.^[47] found that the root of the Surat Thani cultivar of *C. zizanioides* (L.) showed antimicrobial activity due to presence of vetiverin compound in oil. Jayashree et al.^[48] reported that the root extracts of vetiver showed maximum zones of inhibition of 30 mm against *Staphylococcus*

aureus and 33 mm against *Candida albicans*, while the lowest zones of inhibition were 24 mm against *Escherichia coli* (*E. coli*) and 28 mm against *Cryptococcus neoformans*, respectively.

Antibacterial activity:

Numerous studies have revealed that vetiver EO has antibacterial properties that can combat a variety of bacterial strains, including *Staphylococcus aureus*, *Streptococcus pyogenes*, *E. coli*, *Corynebacterium ovis*, *Mycobacterium smegmatis*, *Acinetobacter baumannii*, *Aeromonas veronii*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*^[49,1,50,43]. Nantachit et al.^[47] and Pawar et al.^[50] found that crude root extract from the Surat Thani cultivar of *V. zizanioides* (L.) Nash exhibited antibacterial efficacy against four pathogenic microorganisms. The EO of *C. zizanioides* contains alkaloids, terpenoids, flavonoids, saponins, phenols, and tannins that, either singly or in combination, have antibacterial characteristics^[51]. Putiyanan et al.^[52] showed that *C. zizanioides* (L.) has antibacterial action against *Pseudomonas aeruginosa*, *E. coli*, and *Staphylococcus aureus*. conducted an investigation and proposed that cedr-8-en-13-ol, 6-isopropenyl-4, 8a-dimethyl-1,2,3,5,6, 7,8,8a-octahydro-naphthalen-2-ol, δ -Selinene, γ -Gurjunenepoxide-(2), and solavetivone are the most important compounds in the antibacterial activity of vetiver oil against the Multidrug

Resistance (MDR) bacteria models used in their study.

In a study by Efe *et al.*^[25] found that the vetiver EO had strong antibacterial activity for *Enterobacter cloacae*, *Enterococcus faecalis*, *E. coli*, and *Proteus vulgaris* in the Minimum Inhibitory Concentration (MIC) values, 15.63, 31.25, 15.63, and 15.63 g/ml, respectively. Similarly, Oliveira *et al.*^[13] found that when tested against *Prevotella nigrescens*, *Fusobacterium nucleatum*, *Prevotella melaninogenica*, and *Aggregatibacter actinomycetemcomitans*, vetiver oil exhibited strong antibacterial activity, with MIC values ranging from 22-62.5 µg/ml and Minimum Bactericidal Concentration (MBC) values ranging from 22-400 µg/ml.

Antioxidant activity:

Vetiver oil showed antioxidant activity and has applications in both fragrance and medicine. Two separate *in vitro* assays, the 1,1-Diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging assay and the Fe²⁺ metal chelating assay, were used to assess the antioxidant capabilities of vetiver oil^[50-54]. Many studies have shown the biological activity of the EO of *V. zizanioides* roots. Kim *et al.*^[55] found that the antioxidant property of vetiver has been identified by three distinct compounds vetivenene, both β-vetivone and α-vetivone. The ethanolic extract of vetiver root is used for various antioxidant activities, lowering oxidation ability, superoxide anion radical capturing ability, and overall antioxidant capacity^[44]. Subhadra *et al.*^[49] had conducted research and found that *V. zizanioides* scavenges free radicals, reducing the harm caused by oxidative stress in many illness states, and acting as a possible natural source of antioxidant. Similarly, in a study Samman *et al.*^[36] concluded that the overall antioxidant capacity of the vetiver EO was 75.5 % at 0.1 mg/ml (100 ppm), and the total phenolic content was 6.7 mg Gallic Acid Equivalents(GAE)/g.

The antioxidant activity of EO is depending upon their chemical composition particularly the presence of compound which have radicle function. Soidrou *et al.*^[34] reported that antioxidant action is dosage dependant and found that the concentration of 0.5 mg/ml was highest concentration which shows the strong activity (60.43 %). The concentration of 0.03125 mg/ml, the lowest concentration tested, showed the lowest antioxidant activity, 52.74 %.

Anti-inflammatory activity:

The action that relieves pain is anti-inflammatory. Vetiver EO had an anti-inflammatory activity. It is mostly utilised as an anti-inflammatory in the neurological and circulatory systems. It is also effective for treating inflammations caused by sunburn, dehydration, and dry winds^[56,44]. Chou *et al.*^[11] found that 25 compounds that make up *V. zizanioides*-EO, with cedr-8-en-13-ol (12.4 %), α-amorphene (7.80 %), β-vetivone (5.94 %), and α-gurjunene (5.91 %) being the main constituents. Vetiver EO may suppress the inflammatory responses of Lipopolysaccharide (LPS) -stimulated RAW 264.7 macrophages, including nitric oxide production and cell apoptosis, through the regulation of the expression of the inflammation-related enzymes Heme Oxygenase-1 (HO-1), inducible Nitric Oxide Synthase (iNOS) and cyclooxygenase-2 and the inflammatory cytokines Tumor Necrosis Factor (TNF)-α, Interleukin (IL)-1β and Interferons (IFN-β). Lima *et al.*^[33] found that vetiver EO can stop abdominal writhes, reduce the second phase of formalin-induced pain, and inhibit carrageenan-induced inflammation, providing evidence of its analgesic and anti-inflammatory properties.

Anti-cancer activity:

Vetiver EO is cytotoxic to cancer cells^[37]. Chen *et al.*^[57] conducted research and found that vetiver oil reduced the development of SiHa cervical cells by 89 %, CaSki cervical cells by 88 %, and MCF-7 breast cancer cells by 89 %. Similar results were found by Hanifa *et al.*^[58] that the substances found in vetiver oil principally responsible for the cytotoxic actions are β-caryophyllene and α-humulene, which especially influence Triple Negative Breast Cancer (TNBC) directed at cannabinoid receptor 2. South Indian vetiver grown for commercial purposes contained 0.37 % β-caryophyllene (Table 2). Additionally, β-caryophyllene has received a lot of attention as a potential cancer treatment and prevention agent^[58]. Sinha *et al.*^[59] reported that in human lymphocytes, vetiver acetate was found to be considerably genotoxic and cytotoxic at high doses, but vetiver oil is safe for consumption at moderate quantities. In a study Chitra *et al.*^[60] found that the root extract of vetiver showed cytotoxicity when applied to a cancer cell line. The result reveals that the human breast cancer MCF-7 cell line has an half maximal Inhibitory

Concentration (IC_{50}) concentration ranging from 5-50 g/ml. MIC was seen between 31 and 37 g/ml.

Anti-fungal activity:

Phytopathogenic fungi were examined for resistance to the antifungal effects of sesquiterpenoids of vetiver oil^[42,50,1]. The antifungal action of vetiver has also been demonstrated in numerous investigations against *Alternaria alternata*, *Fusarium oxysporium*^[50], *Rhizoctonia bataticola*, *Sclerotium rolfsii*, *Aspergillus niger*, *Candida albicans*, and *Cryptococcus neoformans*^[37,43]. Khusinodiol monobrosylate, one of the several substances examined, has been found to be an efficient antifungal agent against fungi^[1]. Putiyanan *et al.*^[52] found that six different species of vetiver have been found to have antifungal activity against *Trichophyton mentagrophytes*. Dubey *et al.*^[38] studied that antifungal activity against the phytopathogenic fungus *Rhizoctonia solani*, 50 % Effective Dose (ED_{50}) ($\mu\text{g/ml}$) showed that South Indian vetiver oil was substantially more effective than North Indian oil. Similarly, Samman *et al.*^[36] conducted a study and found that vetiver oil showed a considerable inhibition in radial growth of all the studied fungus species, including *Fusarium oxysporum*, *Alternaria citri*, and *Rhizoctonia solani* at the lowest application rate (500 g/ml), and it was strengthened when concentration was raised to (1000 g/ml). In a Study by Powers *et al.*^[61] revealed that EOs from vetiver shown significant antifungal activity against *C. neoformans* (MIC=20 g/mL) and cytotoxic action against MCF-7 cells (IC_{50} =23.9 g/mL). Sesquiterpene alcohols and khusimol are abundant in vetiver EO. The development of *A. niger* and *C. albicans* was inhibited by vetiver oil (MIC=78 and 313 g/ml),

CONCLUSION

The EO of *C. zizanioides* shows promise for usage in novel care products and that it can be further used to develop new medications, cosmetic and various industrial products. The complexity of the oils' chemical makeup and the variety of their constituents and chemical structures are responsible for the wide range of biological activities. It can be concluded that vetiver oil has been shown effectiveness against bacterial, fungal, microbial, cancer diseases with great economic importance. It is also a complex mixture of various sesquiterpenes, hydrocarbons, flavonoids,

saponins, phenols, and tannins, and is used in various medical and cosmetic industries.

Furthermore, as biological activity can strongly vary depending on a species chemical makeup inside it, future research should concentrate on plant culture and EO standardisation. Various chemical compounds of vetiver are still unidentified and needs to identify their properties and uses in various pharmacological activities.

Acknowledgement:

Authors acknowledged the support received from the central library staff members, Guru Ghasidas University for the literature search.

Authors' Contributions:

Ankit Pandey: Data curation, data analysis, drafting, writing, and formatting. S.C. Tiwari: Conceptualization, draft correction and editing, and suggestions for improvements.

Conflict of interests:

The authors declared no conflict of interests.

REFERENCES

1. Chahal KK, Bhardwaj U, Kaushal S, Sandhu AK. Chemical composition and biological properties of *Chrysopogon zizanioides* (L.) Roberty syn. *Vetiveria zizanioides* (L.) Nash-A Review. Indian J Nat Prod Resour 2014;6(4):251-60.
2. Yanto E, Agustian E, Sulastawaty A. Simple purification of vetiver oil by multiglass plate system for quality improvement. Indones J Appl Sci 2017;18(02):124-31.
3. Raviprasad Sajjan M, Venugopal CK. Studies on the effect of planting methods and nutrition on growth, yield and essential oil content in vetiver (*Vetiveria zizanioides* (L.) Nash). Int J Chem Stud 2017;5(3):225-9.
4. Sajjan MR, Venugopal CK, Chandranath HT, Naik BK, Mokashi AN. Physico-chemical properties of essential oil in vetiver (*Vetiveria zizanioides* (L.) Nash) as influenced by different planting methods and nutrition. Int J Chem Stud 2019;7(1):1443-7.
5. Priyadarshani ND, Amarasinghe MK, Subasinghe S, Palihakkara IR, Kumarasinghe HK. Effect of potting mixture on plant growth, yield and oil quality of vetiver (*Vetiveria zizanioides* L). Trop Agric Res and Ext 2011;13(4).
6. Bhushan B, Sharma SK, Singh T, Singh L, Arya H. *Vetiveria zizanioides* (Linn.) Nash: A pharmacological overview. Int Res J Pharm 2013;4(7):18-20.
7. Lunz K, Stappen I. Back to the roots-an overview of the chemical composition and bioactivity of selected root-essential oils. Molecules 2021;26(11):3155.
8. Lavania UC. Primary and secondary centres of origin of vetiver and its dispersion. In International vetiver conference, 2. Proceedings. Thailand: TVN 2000:424-426.
9. Lal RK. On genetic diversity in germplasm of vetiver '*Vetiveria zizanioides* (L.) Nash'. Ind Crop Prod 2013;43:93-8.

10. Singh N, Singh VR, Lal RK, Verma RS, Mishra A, Yadav R. Quantification of genotypic and chemotypic diversity for elite clone selection with high-quality essential oil traits in vetiver [*Chrysopogon zizanioides* (L.) Roberty]. *J Essent Oil-Bear Plants* 2019;22(4):1150-62.
11. Chou ST, Lai CP, Lin CC, Shih Y. Study of the chemical composition, antioxidant activity and anti-inflammatory activity of essential oil from *Vetiveria zizanioides*. *Food Chem* 2012;134(1):262-8.
12. Alifano P, del Giudice L, Tala A, de Stefano M, Maffei ME. Microbes at work in perfumery: the microbial community of vetiver root and its involvement in essential oil biogenesis. *Flavour Fragr J* 2010;25(3):121-2.
13. Oliveira TA, Vieira TM, Esperandim VR, Martins CH, Magalhães LG, Miranda ML, *et al.* Antibacterial, antiparasitic, and cytotoxic activities of chemical characterized essential oil of *Chrysopogon zizanioides* roots. *Pharmaceuticals* 2022;15(8):967.
14. David A, Wang F, Sun X, Li H, Lin J, Li P, *et al.* Chemical composition, antioxidant, and antimicrobial activities of *Vetiveria zizanioides* (L.) nash essential oil extracted by carbon dioxide expanded ethanol. *Molecules* 2019;24(10):1897.
15. Amalia N, Sukrasno S, Fidrianny I. Isolation of khusimol from the root of vetiver (*Vetiveria zizanioides* L. Nash) grown in samarang-garut and the study of its profile after harvesting. *Lett Appl Nano Biosci* 2022;11(2):3549-59.
16. Mallavarapu GR, Syamasundar KV, Ramesh S, Rao BR. Constituents of South Indian vetiver oils. *Nat Prod Commun* 2012;7(2):223-225.
17. Kusuma HS, Rohadi TI, Daniswara EF, Altway A, Mahfud M. Preliminary study: Comparison of kinetic models of oil extraction from vetiver (*Vetiveria zizanioides*) by microwave hydrodistillation. *Korean Chem Eng Res* 2017;55(4):574-7.
18. Raja MB, Rajamani K, Suresh J, Joel AJ, Uma D. Chemical composition of vetiver root oil obtained by using GCMS analysis. *J Pharmacogn Phytochem* 2018;7(6):1709-13.
19. Lavania UC. Other uses and utilization of vetiver: Vetiver oil. In the third International vetiver conference, Guangzhou, China 2003:475.
20. Lavania UC. Vetiver in India: Historical perspective and prospective for development of specific genotypes for environmental or industrial application. *J Agric Food Environ Sci* 2008:40-7.
21. Feng Z. Vetiver system technology: Application and development in china from 1998 to 2015.
22. Chowdhury AR, Kumar D, Lohani H. GC-MS analysis of essential oils of *Vetiveria zizanioides* (Linn.) nash roots. *Fafai J* 2002;4:33-5.
23. Peng HY, Lai CC, Lin CC, Chou ST. Effect of *Vetiveria zizanioides* essential oil on melanogenesis in melanoma cells: Downregulation of tyrosinase expression and suppression of oxidative stress. *Sci World J* 2014;2014(1):213013.
24. Bhuiyan MN, Chowdhury JU, Begum J. Essential oil in roots of *Vetiveria zizanioides* (L.) nash ex small from Bangladesh. *Bangladesh J Bot* 2008;37(2):213-5.
25. Efe D. The evaluation of the antibacterial activity of *Vetiveria zizanioides* (L.) nash grown in Giresun. *Alinteri J Agric Sci* 2019;34(1):21-4.
26. Kirici S, Inan M, Turk M, Giray ES. To study of essential oil and agricultural properties of vetiver (*Vetiveria zizanioides*) in the Southeastern of Mediterranean. *Adv Environ Biol* 2011:447-52.
27. Danh LT, Mammucari R, Truong P, Foster N. Response surface method applied to supercritical carbon dioxide extraction of *Vetiveria zizanioides* essential oil. *Chem Eng J* 2009;155(3):617-26.
28. Pripdeevech P, Wongpornchai S, Marriott PJ. Comprehensive two-dimensional gas chromatography-mass spectrometry analysis of volatile constituents in Thai vetiver root oils obtained by using different extraction methods. *Phytochem Anal* 2010;21(2):163-73.
29. Pripdeevech P, Wongpornchai S, Promsiri A. Highly volatile constituents of *Vetiveria zizanioides* roots grown under different cultivation conditions. *Molecules* 2006 25;11(10):817-26.
30. Champagnat P, Figueredo G, Chalchat JC, Carnat AP, Bessiere JM. A study on the composition of commercial *Vetiveria zizanioides* oils from different geographical origins. *J Essent Oil Res* 2006;18(4):416-22.
31. Hammam K, Amer A, Noreldin T. Vetiver (*Vetiveria zizanioides* L.) yield and its water use efficiency affected by different plant populations under reclaimed soil conditions. *J Med Plant Res* 2019;7(5):126-34.
32. Thubthimthed S, Thisayakorn K, Rerk-am U, Tangstirapakdee S, Suntornatanasat T. Vetiver oil and its sedative effect. In the 3rd International vetiver conference, Guangzhou, China 2003:492-4.
33. Lima GM, Quintans-Júnior LJ, Thomazzi SM, Almeida EM, Melo MS, Serafini MR, *et al.* Phytochemical screening, antinociceptive and anti-inflammatory activities of *Chrysopogon zizanioides* essential oil. *Braz J Pharmacogn* 2012;22:443-50.
34. Soidrou SH, Farah A, Satrani B, Ghanmi M, Lachkar M, Mohamed AS. Chemical composition, antioxidant and antimicrobial activity of *Vetiveria zizanioides* roots essential oil harvested in Ndzuwani, Comoros. *World J Pharm Pharm Sci* 2020;1(10):82-95.
35. Tunsaringkarn SI, Rungsiyothin A, Ruangrunsi NV. Pharmacognostic specification of *Vetiveria zizanioides* roots in Thailand. *J Health Res* 2008;22(1):9-14.
36. Samaan M, Ebid M, Thabet M. GC-MS analysis, antioxidant capacity and antimicrobial action of *Vetiveria zizanioides*. Essential oil cultivated in North Egypt. *J Plant Prod* 2022;13(9):699-704.
37. Suyono H, Susanti D. The medical benefits of vetiver essential oil. In proceedings of the 2nd International conference of essential oils. 2020;1:9-12.
38. Dubey N, Raghav CS, Gupta RL, Chhonkar SS. Chemical composition and antifungal activity of vetiver oil of North and South India against *Rhizoctonia solani*. *J Nat Pestic Res* 2010;22(1):63-7.
39. Kadarohman A, Dwiyaniti G, Kadarusman E. Quality and chemical composition of organic and non-organic vetiver oil. *Indones J Chem* 2014;14(1):43-50.
40. Putrawan ID, Farda E. Simple vacuum distillation of vetiver oil from smallholders for quality improvement. *Glob J Res Eng Chem* 2015;15(3).
41. Khalil M, Ayoub S. Analysis of the essential Oil of *Vetiveria nigritana* (Benth) Stapf root growing in Sudan. *J Med Plant Res* 2011;32(5):7006-10.
42. Saraswathi KT, Jayalakshmi NR, Vyshali P, Kameshwari

- MS. Comparative study on essential oil in natural and *in vitro* regenerated plants of *Vetiveria zizanioides* (Linn.) Nash. Am-Eurasian J Agric Environ Sci 2011;10(3):459-63.
43. Efe D, Karakose M, Karacelik AA, Ertan B, Seker ME. GC-MS analyses and bioactivities of essential oil obtained from the roots of *Chrysopogon zizanioides* (L.) Roberty cultivated in Giresun, Turkey. Turk J Chem 2021;45(5):1543-50.
 44. Zahoor S, Shahid S, Fatima U. Review of pharmacological activities of *Vetiveria zizanioides* (Linn) Nash. J Basic Appl Sci 2018;14:235-8.
 45. Sangeetha D, Stella D. Screening of antimicrobial activity of vetiver extracts against certain pathogenic microorganisms. Int J Pharm Biol Arch 2012;3:197-203.
 46. Soni A, Dahiya P. Screening of phytochemicals and antimicrobial potential of extracts of *Vetiveria zizanioides* and *Phragmites karka* against clinical isolates. Int J Appl Pharm 2015:22-4.
 47. Nantachit K, Bunchoo M, Khantava B, Khamvan C. Antimicrobial activity of alkaloid from roots of *Vetiveria zizanioides* (L.) Nash ex Small. Thai Pharm Health Sci J 2010;5(2):99-102.
 48. Jayashree S, Rathinamala J, Lakshmanaperumalsamy P. Antimicrobial activity of *Vetiveria zizanioides* against some pathogenic bacteria and fungi. Medic Plants Int J Phytomed Relat Ind 2011;3(2):151-6.
 49. Varadharajan S, Kuppusamy A, Muthuswamy U, Andichettiarhirumalasia S, Rajakannu S. *In vitro* antioxidant activity of *Vetiveria zizanioides* root extract. Tanzan J Health Res 2010;12(4):265-71.
 50. Pawar SH, Patil AB, Prasad DN. Development, and analysis of *Vetiveria zizanioides* for effectiveness as an herbal mosquito repellent. Int J Food Nutr 2012;11(11):383-94.
 51. Kumar SS, Gayathri K. Chemical characterization of *Vetiveria zizanioides* Linn root. Int J Pharm Bio Sci 2016;7(4):689-95.
 52. Putiyanan S, Nantachit K, Bunchoo M, Khantava B, Khamwan C. Pharmacognostic identification and antimicrobial activity of *Vetiveria zizanioides* (L.) Nash root. Chiang Mai Med Bull 2005;44(3):85-90.
 53. Ramírez-Rueda RY, Marinho J, Salvador MJ. Bioguided identification of antimicrobial compounds from *Chrysopogon zizanioides* (L.) Roberty root essential oil. Future Microbiol 2019;14(14):1179-89.
 54. Grover M, Behl T, Bungau S, Aleya L. Potential therapeutic effect of *Chrysopogon zizanioides* (Vetiver) as an anti-inflammatory agent. Environ Sci Pollut Res Int 2021;28:15597-606.
 55. Kim HJ, Chen F, Wang X, Chung HY, Jin Z. Evaluation of antioxidant activity of vetiver (*Vetiveria zizanioides* L.) oil and identification of its antioxidant constituents. J Agric Food Chem 2005;53(20):7691-5.
 56. Balasankar D, Vanilarasu K, Preetha PS, Rajeswari S, Umadevi M, Bhowmik D. Traditional and medicinal uses of vetiver. J Med Plants Stud 2013;1(3):191-200.
 57. Chen F, Wang X, Kim HJ. Antioxidant, anticarcinogenic and termiticidal activities of vetiver oil. In proceeding of third international Vetiver conference, Guangzhou, China 2003.
 58. Hanifa M, Wulandari R, Zulfin UM, Nugroho EP, Haryanti S, Meiyanto E. Different cytotoxic effects of vetiver oil on three types of cancer cells, mainly targeting CNR2 on TNBC. Asian Pac J Cancer Prev 2022;23(1):241.
 59. Sinha S, Jothiramajayam M, Ghosh M, Mukherjee A. Evaluation of toxicity of essential oils palmarosa, citronella, lemongrass and vetiver in human lymphocytes. Food Chem Toxicol 2014;68:71-7.
 60. Chitra T, Jayashree S, Rathinamala J. Evaluation of anticancer activity of *Vetiveria zizanioides* against human breast cancer cell line. Int J Pharm Sci Res 2014;6:164-6.
 61. Powers CN, Osier JL, McFeeters RL, Brazell CB, Olsen EL, Moriarity DM, *et al.* Antifungal and cytotoxic activities of sixty commercially-available essential oils. Molecules 2018;23(7):1549.