

Chemical Constituents and Biological Activities of Genus *Picrorhiza*: An Update

V. PRAKASH*, ANJANA KUMARI, HARPREET KAUR, M. KUMAR, S. GUPTA¹ AND R. BALA²

Department of Chemistry, M.M. University, Sadopur-Ambala, Haryana-134 007, ¹M. M. College of Pharmacy, M. M. University, Mullana, Haryana-133 203, ²Department of Chemistry, Guru Nanak Dev University, Amritsar-143 005, India

Prakash *et al.*: Chemical Constituents and Biological Activities of Genus *Picrorhiza*

This review article on *Picrorhiza* genus comprehensively summarized the characterization, morphology, distribution, traditional uses, chemical constituents and biological activities of isolated individual constituents as well as plant extracts. It belongs to the Scrophulariaceae family, a small genera with two species, which are mainly found in Western and Eastern Himalayas at an altitude of 3000-5200 m. It has been used traditionally to cure many diseases like asthma, arthritis, cancer, diabetes, diarrhea, dyspepsia, fever, gastrointestinal problems, jaundice, urinary disorders. Its phytochemical analysis reported the presence of glycosides, aromatic esters, bis-iridoid, phenyl propenoids, alcoholic compounds and fatty acids. Chemical constituents of the genus *Picrorhiza* and its crude extracts with various solvents exhibit biological activities such as antimicrobial, anticancer, antiasthmatic, antidiabetic effects.

Key words: *Picrorhiza*, glycosides, antimicrobial, anticancer, phytochemicals

Plants and their parts have been used for medicinal uses since immemorial time. The first evidence marked on clay tablets in the cuneiform script is from Mesopotamia 2600 BC where the recipes of drug preparation have been written^[1-3]. It is a universally accepted principle that human beings and diseases are born simultaneously. Vedas^[4], Quran^[5-7], Buddhism^[8] and Bible^[9] are the religious texts that have described diseases, drugs, and therapies. The entire knowledge of *Ayurveda* is derived from *Vedic* literature. *Vedic* system of medicine is the root of *Ayurveda* and it is supposed to be the science of all aspects of life^[10]. Ancient philosophy deliberates the “secret of life and beyond” for health and medicinal plants^[11]. No plant of this earth is useless from a medicinal point of view^[12].

The Himalayas, a mountain range in Asia covering mainly India, Pakistan, China, Tibet, and Nepal is known as a treasure house of the medicinal plants with around 8000 species in which 1748 species are known to have medicinal value^[13-16]. A majority of the rural population (~90 %) depends upon the medicinal plants as a source of remedies^[17-21]. Alone in India about 2000 tons of herb consumed annually^[22-26]. Globally, the excessive use of synthetic drugs and their side effects have attracted the researchers towards herbals medicine in past years. Plants contain many different phytochemical

compounds that are individually effective enough to treat chronic and infectious diseases^[27,28].

The genus *Picrorhiza* is well known for its medicinal values since the ancient as well as modern period. There are not many reviews on *Picrorhiza*, except a few such as cellular differentiation, regeneration and secondary metabolite production in medicinal *Picrorhiza* species^[29], reviews of its chemical constituents^[30], phytopharmacological review on genus *Picrorhiza*^[31], *Picrorhiza kurroa*: an ethnopharmacologically important plant species of the Himalayan region^[22]. Therefore, there is a need to prepare an updated detailed review on the genus *Picrorhiza*. This review mainly deals with the isolation of plant extracts and individual chemical constituents of *Picrorhiza* and studies of biological activities shown by an individual chemical constituent as well as on solvent extracts carried out so far in the last decade, based upon the literature survey through PubMed, Mendeley, Ex Prisma.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms

*Address for correspondence
E-mail: vinitprakash@gmail.com

Classification:

Picrorhiza is commonly called *karu*^[32], while scientifically it is placed in Kingdom: Plantae; Class: Dicotyledon; Order: Scrophulariales; Family: Scrophulariaceae; Genus: *Picrorhiza*; Species: *kurroa*, *scrophulariiflora*; Botanical name: *P. kurroa* Royal ex Benth and *P. scrophulariiflora* Pennel^[31].

Morphology:

The genera *Picrorhiza* is derived from 'picros' and 'rhiza', which means bitter root^[33-35]. Genus *Picrorhiza* has two species namely *Picrorhiza kurroa* and *Picrorhiza scrophulariiflora*^[36]. *P. kurroa* is smaller in height (5-9 cm) with smaller leaves (7-11 cm). The leaves of the plant are flat, oval, saw-toothed, five narrowed to a winged stalk, and sharply serrated. The flowers are white or pale purple and borne on a tall spike and its corolla is much smaller (0.8 cm), five-lobed to the middle, and with very much longer stamens. The rhizome is 2.5-12 cm long, sub-cylindrical, straight, or to some extent curved, and the external surface is coarse^[37-40].

P. scrophulariiflora is longer than *P. kurroa* in height (8-12 cm) with longer leaves. The leaves of the plant are flat, oval, saw-toothed and its leaf blades are 10-14 cm in length. The flowers are dark blue-purple in color arising from a rosette of conspicuously serrated leaves and its corolla is 1.5 cm. Rhizomes are 2.5-12 cm long, sub-cylindrical, straight, or to somewhat curved and the rhizomes are evanescent inside^[37].

Distribution:

P. kurroa occurs mainly in the Western Himalayas at an altitude of 3000-5000 m^[34,41-44], while *P. scrophulariiflora* is found mainly in the Eastern Himalayas at an altitude of 4300-5200 m. *P. scrophulariiflora* species of this genus is found only in Nepal^[45,46].

Traditional uses:

The World Health Organization (WHO) defined that the traditional medicine is a total sum of all knowledge and practices, which is explainable or not and it is used in diagnosis, prevention and elimination of physical, mental, or social imbalance and relying exclusively on practical experience and observation handed down from generation to generation, whether verbally or in writing^[46]. Traditional knowledge is passing from generation to generation without the aid of any documentation or keeping written records. This knowledge will be lost with succeeding generations if

it is not rapidly researched and recorded^[18]. Therefore, its documentation is of paramount importance and urgent so that it can be preserved and conserved^[47,48]. Ethnobotanical surveys are effective methods in documenting and identifying medicinal plants used in the traditional knowledge system^[49]. Both the species *P. kurroa* and *P. scrophulariiflora* have been used traditionally for the treatment of a large number of diseases long before the time^[50,51]. The different parts of *P. kurroa* have been used to cure various diseases. The rhizomes of *P. kurroa* are used to cure asthma, arthritis, cancer, diabetes, dyspepsia, fever, gastrointestinal problems, jaundice, leukoderma, piles, snake bite, and urinary disorders.

The seeds of *P. kurroa* are used to treat diarrhea, fever, liver, and upper respiratory tract disorders, its leaves are used to cure diseases such as bilious fever and chronic infections, and its flowers are also used for the treatment of fever and blood purification^[41,42,52-77].

Traditionally, stems of *P. scrophulariiflora* have been used to cure arthritis, asthma, choleric, jaundice, liver disorder while its roots are useful for the treatment of bacterial infections, fever, jaundice, liver disorder, and inflammatory diseases^[78-81]. *P. scrophulariiflora* is also used in traditional Chinese medicine for treating dysentery, jaundice and rheumatism^[82].

Isolation of chemical constituents:

During the said period of review around 53 compounds have been isolated from these 2 species of genus *Picrorhiza*, out of which 38 are glycosides, 6 aromatic esters, 4 bis-iridoids, 4 phenylpropanoids and an alcoholic compound. From *P. kurroa*, a total of 36 compounds have been isolated, which included 26 glycosides, 6 aromatic esters and 4 bis-iridoids from different parts of the plants whereas 17 compounds have been isolated from *P. scrophulariiflora*, which contained only 12 glycosides, 4 phenylpropanoids and an alcoholic compound (Table 1). The structures of these compounds are given in fig. 1 (excluding compounds reported by Sah and Varshney^[30]).

Glycosides are formed from the combination of a hydroxyl group of a simple sugar with another compound. From the genus *Picrorhiza*, 8 types of glycosides have been isolated, which included iridoid glycosides (23), aromatic aldehyde glycoside (1), aromatic aldehyde diglycoside (1), diphenolic tetra glycoside (1), phenolic glycosides (5), phenylpropanoid glycoside (2), secoiridoid glycosides (2) and hydroquinone glycosides (3) from different parts of the

TABLE 1: CHEMICAL CONSTITUENTS AND BIOLOGICAL ACTIVITIES OF GENUS *PICRORHIZA*

S. No.	Chemical constituents (No.)	Plant species	Plant Part	Solvent employed	Biological activities	Reference
Glycosides						
1.	[6-[(1a,1b,2,5a,6,6a)-6-hydroxy-1a-(hydroxymethyl)-2,5a,6,6a-tetrahydro-1bHoxireno[5,6]cyclopenta [1,3-c]pyran-2-yl]oxy]-3,4,5-trihydroxyoxan-2-yl] methyl-3 phenylprop-2-enoat. (1) IG	P.K.	Rhizomes; Rhizomes and leaves	Methanol; Methanol: water (1:1); Ethanol and Water	Anticancer activity; Antiarthritis activity; Antidiabetic activity; Hepatoprotective activity.	[75,84,86,88, 89,90,92]
2.	[2-(hydroxymethyl)-10-[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxy-3,9-dioxatricyclo[4.4.0.0 ^{2,4}]dec-7-en-5-yl] 4-hydroxy-3-methoxybenzoate. (2) IG	P.K.	Rhizomes; Rhizomes and leaves	Methanol; Methanol: water; Methanol and Water; Ethanol and Water	Antiarthritis activity; Antidiabetic activity; Hepatoprotective activity.	[75,84,88, 89,90,92]
3.	[5-hydroxy-10-[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxy-3,9-dioxatricyclo[4.4.0.0 ^{2,4}]dec-7-en-2-yl] methyl 4-hydroxy-3-methoxybenzoate. (3) IG	P.K.	Rhizomes	Ethanol; Methanol	Antimicrobial activity; Anticancer activity.	[85,86]
4.	[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl] 3-methoxy-4-[3-phenylprop-2-enoyl]oxybenzoatedihydrate. (4) IG	P.K.	Rhizomes	Water; Methanol	Antidiabetic activity; Antimicrobial activity	[86,87]
5.	1-(4-hydroxy-3-methoxyphenyl)ethanone. (5) IG	P.K.	Rhizomes	Ethanol and Water	Antidiabetic activity.	[90]
6.	Vanillin- α -D-glucopyranoside. (6) AAG	P.K.	Rhizomes	Methanol	-----	[91]
7.	Picaldehyde 4-O- α -D-glucopyranosyl-(6'→1'')- O- α -D-glucopyranoside. (7) AADG	P.K.	Rhizomes	Methanol	-----	[91]
8.	Picrorhizaoside-A. (8) IG					
9.	Picrorhizaoside-B. (9) IG					
10.	Picrorhizaoside-C. (10) IG					
11.	Picrorhizaoside-D. (11) IG	P.K.	Rhizomes	Methanol	Hyaluronidase inhibitory activity	[110]
12.	Picrorhizaoside-E. (12) IG					
13.	Picrorhizaoside-F. (13) IG					
14.	Picrorhizaoside-G. (14) IG					
15.	3-methoxy-4-hydroxyphenyl- <i>n</i> -butanyl- α -O-D-glucopyranosyl-(6a→1b)- α -O-D-glucopyranosyl-(6b→1c)- α -O-D-glucopyranosyl-(6c→1d)- α -O-D-glucopyranosyl-4d-	P.K.	Rhizomes	Methanol	-----	[91]
16.	3'-methoxy-4'-hydroxyphenyl- <i>n</i> -pent-7',9'-dien-11'-oate. (15) DPTG					
17.	3-phenylprop-2-enoic acid. (16) IG					
18.	3-(4-hydroxy-3-methoxyphenyl)prop-2-enoic acid. (17) IG					
19.	6-Hydroxy-1a-(hydroxymethyl)-1a,1b,2,5a,6,6a-hexahydrooxireno[4,5]cyclopenta[1,2-c]pyran-2-yl 6-O-[(2E)-3-(4-hydroxy-3-methoxyphenyl)-2-propenoyl]- β -D-glucopyranoside. (18)IG	P.K.	Rhizomes	Hydro-alcoholic	-----	[92]
19.	Scrophuloside. (19) IG					
20.	Veronicoside. (20) IG					


21.	Abeloside A. (21) IG					
22.	Abeloside B. (22) IG					
23.	Sylvestroside IV dimethyl acetal. (23) IG	P.K.	Stems	n- Butanol	-----	[93]
24.	Sweroside. (24) IG					
25.	8-Epi-loganin. (25) IG					
26.	8-Epi-loganic acid. (26) IG					
27.	2-(3,4-dihydroxyphenyl)-3,4-dihydro-2H-chromene-3,5,7-triol. PG					
28.	2-(3,4-dihydroxyphenyl)-5,7-dihydroxychromen-4-one. PG	P.S.	Rhizomes	95% Ethanol	-----	[30,97]
29.	Scroside G. PG					
30.	7-Hydroxy-coumarine. (27)PG					[113]
31.	Cinnamic acid. (28) PG	P.S.	Rhizomes	n-Butanol	-----	
32.	Scroside H. (29)PPG					
33.	Scroside I. (30)PPG	P.S.	Roots	Ethanol	-----	[114]
34.	picrogentioside II. (31)SIG	P.S.	Roots	Ethanol	-----	[98]
35.	picrogentioside D. (32)SIG	P.S.	Roots	Methanol	-----	[115]
36.	Hebitol III. (33)HQQ	P.S.		Methanol		
37.	Scrophuloside C. (34)HQQ		Roots		-----	[116]
38.	Scrophuloside D. (35)HQQ					
Aromatic Ester						
39.	3- Methoxy-4-decanoxy benzoic acid. (36)					
40.	3-Methoxy-4-tetradecanoxy-phenyl n-pent-7,9-diene-11-al. (37)	P.K.	Rhizomes	Macerated with aqueous	-----	[100]
41.	3- Methoxy-4-dodecanoxy- phenyl-n-pent-7,9-dien-11-al. (38)					
42.	3- Methoxy-4-dodecanoxy phenyl- n-pent-7,9-dien-11-al. (39)					
43.	3-Methoxy-4-tetradecanoxy-phenyl n-pent-7,9-diene-11-al. (40)	P.K.	Rhizomes	Methanol	-----	[91]
44.	3-Methoxy-4-decanoxy benzoic acid. (41)					
Bis-iridoid						
45.	Saungmaygaoside A. (42)					
46.	Saungmaygaoside B. (43)					
47.	Saungmaygaoside C. (44)	P.K.	Stems	n- Butanol	-----	[93]
48.	Saungmaygaoside D. (45)					
Phenylpropenoids						
49.	Luteolin -7-O-β-D-glucoside.					
50.	Gallic acid.	P.S.				
51.	Isoferulic acid.		Rhizomes	95% Ethanol	-----	[30,97]
52.	Vanillic acid					
Alcoholic compounds						
53.	Hexacosanol.	P.S.	Rhizomes	95% Ethanol	-----	[30,97]

P.K.-*Picrorhizakurroa*, P.S.- *Picrorhizascrophulariiflora*, IG- Iridoid glycoside, AAG-Aromatic aldehyde glycoside, AADG- Aromatic aldehyde diglycoside, DPTG-Diphenolic tetra glycoside, PG- Phenolic glycoside, PPG-Phenylpropenoid glycoside, SIG- Secoiridoid glycoside, HQG-Hydroquinone glycoside


Picrorhiza plant. In *P. kurroa* there were 26 glycoside compounds (1-26), reported from the different parts of a plant like rhizomes, leaves and stems using different solvents like methanol, ethanol, n-butanol, chloroform, water, and methanol:water (1:1)^[75,83-93] while in *P. scrophulariiflora* there are 12 glycosides (27-35),

which have been reported from the different parts viz. roots, leaves and stems using different solvents like ethanol, butanol, (95 %) and ethyl acetate^[29,30,83,91,94-99].


In aromatic esters, ester linkage is bonded directly to an aromatic system. Only from the roots of *P. kurroa* species 6 (36-41) aromatic ester have been isolated with




(1) (2) (3) (4) (5) (6)



(7) (8) (9) (10) (11)








Fig. 1: Structures of isolated chemical compounds from genus *Picrorhiza*

water and methanol^[91,100]. In case of bis-iriodids, 2 units are connected by a 1,3-dioxane group. Four bis-iriodids (42-45) have been isolated from the stems of *P. kurroa* with n-butanol^[93]. From the roots of *P. scrophulariiflora* four phenylproponoid compounds have been isolated using (95 %) ethanol as a solvent^[30,97].

Alcohols are organic compounds that contain at least one hydroxyl functional group bound to their aliphatic or aromatic substructure. One alcoholic compound has been isolated from the roots using ethanol as a solvent^[30,97]. In the genus, *Picrorhiza* main phytochemical constituents are glycosides, which have been mainly isolated from rhizomes. The chemistry of glycosides from leaves, seeds, fruits and stems need to be explored.

Phytochemical analysis:

Many researchers used different analytical techniques like UV/Vis, IR, GC-MS, GC-FID, LC-MS, NMR, HPLC, HPTLC, UPLC, for the identification and characterization of new compounds from the genus *Picrorhiza*. A brief summary of *Picrorhiza* species using different analytical techniques are shown in Table 2.

Biological activities:

A wide range of biological activities of isolated chemical constituents and plant extracts, like antimicrobial^[85], hepatoprotective^[92], antioxidant activity^[101], anticancer^[86,102], antiarthritic^[88], antidiabetic^[87],

TABLE 2: PHYTOCHEMICAL ANALYSIS ON GENUS *PICRORHIZA*

Sr. no.	Plant species	Part of plant	Solvent used for extraction	Analytical Technique used	Result	References
1.	<i>P. kurroa</i>	Rhizomes	Methanol	HPTLC	Yield of extract is 40.63% (Rf 0.24)	[117]
2.	<i>P. kurroa</i>	Rhizomes	Methanol	HPTLC	Picroside-1 (Rf 0.51) and kutkoside (Rf 0.63)	[118]
3.	<i>P. kurroa</i>	Rhizomes	Ethanol:water	HPLC-UV	Picroside-1: 3.30%(w/w) Picroside-2: 4.90%(w/w)	[119]
4.	<i>P. kurroa</i>	Rhizomes	Diethyl ether	LC-ESI-MS/MS	cinnamic acid, ferulic acid, kuttoside, picroside-I, picroside-II, picroside-III, scrophuloside-A, and veronicoside.	[92]
5.	<i>P. kurroa</i>	Rhizomes	Ethanol:water	HPLC	Picroside-1: 9.57%(w/w) Picroside-2: 4.30%(w/w)	[90]
6.	<i>P. kurroa</i>	Rhizomes and leaves	Methanol:water	HPLC	Picroside-1: 98.3%(w/w) Picroside-2: 101.3%(w/w)	[89]
7.	<i>P. kurroa</i>	Roots	70% Ethanol	LC-ESI-MS/MS	Picroside III Apocynin, Pikuroside, Picroside IV, Vanillic acid, Picroside II, Kutkoside, Picroside I	[120]
8.	<i>P. kurroa</i>	Rhizomes	Ethanol	ILUAE	picroside I: 2.84% picroside II: 3.57% 6-O-E-feruloyl catalpol:2.20%	[121]
9.	<i>P. kurroa</i>	Rhizomes	Ethanol: water (mother extract) further fractionated using hexane, n-butanol, acetone, methanol and water	HPTLC	Mother extract: (25.6%w/w), hexane (9%w/w), DCM (31%w/w), n-butanol (23%w/w), acetone (11% w/w), methanol (16%w/w), and water (7%w/w). Cucurbitacin B Cucurbitacin D Cucurbitacin E Betulinic acid Picroside 2 Apocynin	[102]
10.	<i>P. scrophulariiflora</i>	Fruits	Ethanol	UPLC	Apocynin, Androsin, Picroside-1, Picroside-2, Picroside-3	[122]
11.	<i>P. kurroa and P. scrophulariiflora</i>	Roots	Methanol	HPTLC	Comparing Picroside-1 and Picroside-2 from two different species of picrorhiza Picroside-1(kurroa): 1.258% Picroside-1 (scrophulariiflora): 1.611% Picroside-2(kurroa): 0.481% Picroside-2 (scrophulariiflora): 0.613%	[123]

immunomodulating^[103,104] have been reported either from the whole plant or specific homogenized parts to prove its medicinal importance (Table 3 and 4).

Iridoid glycosides (3-4) isolated from the rhizomes of *P. kurroa* have shown good antimicrobial activity (cup-plate method) against Gram-positive bacteria, *Bacillus subtilis*, *Staphylococcus aureus*, *Micrococcus luteus*, Gram-negative bacteria, *Pseudomonas aeruginosa*, *Escherichia coli* and fungal strains such as *Aspergillus niger*, *Candida albicans* and *Malasseiza furfur*^[85,87].

The extract prepared using ethanol from the leaves of *P. kurroa* shown antimicrobial activity against the strains of *Staphylococcus*, *Pseudomonas*, and *E. coli* using the agar well diffusion method, and antifungal activity against the fungal strain viz *Candida albicans*, *Candida tropicalis*, *Trichophyton rubrum*, and *Penicillium marneffii*^[105].

Iridoid glycoside (5) extracted from rhizome inhibited tumor invasion and migration of MCF-7 cells^[86]. The rhizome extract of *Picrorhiza* has shown cytotoxicity in XTT assay in MDA-MB-435S (human breast carcinoma), Hep3B (human hepatocellular carcinoma), and PC-3 (human prostate cancer) cell lines^[101]. Mallick *et al.*^[102] studied the anticancer activity of the hydroalcoholic extract of *P. kurroa* and its fractions in breast cancer and cervix cancer cell lines. Oral administration of dichloromethane fraction significantly reduced tumor volume, tumor weight, and percent packed cell volume at a dose of 50 mg/kg, as compared with toxic control group.

Iridoid glycosides (5) extracted with water from the rhizomes of *P. kurroa* have shown antidiabetic activity. Compound (1) was orally administered to streptozotocin-induced diabetic rats at the doses of 100 and 200 mg/kg/day for 14 consecutive days. Plasma insulin levels were measured and the pancreas of the rat was subjected to histopathological investigations. There was evidence of regeneration of β -cells of pancreatic islets of picrorhiza extract-treated group in histopathological examinations. It increased the insulin-mediated translocation of glucose transporter from the cytosol to plasma membrane or increased glucose transporter type-4 appearance, which in turn facilitated glucose uptake by skeletal muscles in diabetic rats^[87]. Aqueous ethanol extract of *P. kurroa* rhizomes demonstrated anti-diabetic activity. Treatment with *P. kurroa* rhizomes extract (100 and 200 mg/kg) for 30 d significantly decreased high blood glucose and restored the normal levels of serum biochemistry^[90].

Saponin and flavonoids have been isolated from ethanol extract of root of this plant, which exhibited antiasthmatic activity in bronchoconstriction and induced airway obstruction method. These saponins have been shown to possess mast cell stabilizing while flavonoids have been reported to possess smooth muscle relaxant and bronchodilator activity^[106].

The ethanol extract of *P. kurroa* leaves were tested on a rat model of anemia induced by intraperitoneal injection of phenylhydrazine at 40 mg/kg for 2 d. The extract was given orally to the rats previously treated with phenylhydrazine at 100 and 200 mg/kg/d, resulted in the increased concentration of hemoglobin, red blood cell number, hematocrit, and reticulocytes rate^[107].

Methanol extract of rhizomes of *P. kurroa* showed antioxidant activity in the superoxide anion radical scavenging method^[108]. Antioxidant activity of the methanol extract was carried out using the radical scavenging assays (DPPH and OH), ferric reducing antioxidant property and thiobarbituric acid assay for testing inhibition of lipid peroxidation^[101].

Kumar *et al.* 2016 studied the hydroalcoholic extract of rhizomes of *P. kurroa*, which showed antiarthritic activity. Additionally, *P. kurroa* rhizome extract significantly inhibited the expression of degrading enzymes, matrix metalloproteinases-3, and matrix metalloproteinases-9 in arthritic rats^[88]. The methanol root extract of *P. kurroa* showed antitrypanosomal activity. It was carried out on a Vero cells and there was a complete kill of trypanosomes^[109]. Picrorhizaosides D (4, IC₅₀ 43.4 μ M) and E (5, 35.8 μ M) have shown strong hyaluronidase inhibitory activity^[110].

Hydroalcoholic extract of rhizomes of *P. kurroa* demonstrated hepatoprotective activity at the dose levels of 50 and 100 mg/kg exemplified by significant restoration of liver enzyme levels and the antioxidant enzymes close control. *Picrorhiza kurroa* hydroalcoholic extract has ameliorative activity from tissue damage against induced hepatotoxicity^[92].

Scrocaffeside A, a compound extracted from rhizomes of *P. scrophulariiflora* has shown immunomodulatory activity against spleen cell suspension of mice. Scrocaffeside A between 5 and 125 μ g/ml stimulated proliferation of splenocytes and their response to polyclonal T cell mitogen concanavalin A and lipopolysaccharide^[111].

Water extract of dried powdered rhizomes of *P. scrophulariiflora*, was given orally in doses of 20, 40, and 80 mg/kg/d for 4 w to rats fed with high-

TABLE 3: BIOLOGICAL ACTIVITIES OF ISOLATED CHEMICAL CONSTITUENTS

Solvent used	Chemical constituents	Strains/Test Animals	Doses and Control	Result	Reference
<i>Picrorhiza kurroa</i>					
Antimicrobial Activity					
Ethanol	[5-hydroxy-10-[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxy-3,9-dioxatricyclo[4.4.0.0 ^{2,4}]dec-7-en-2-yl]methyl 4-hydroxy-3-methoxybenzoate. (3)	<i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> and <i>Escherichia coli</i>	5 mg, 10 mg, 15 mg, 20 mg/75 µl and ciprofloxacin (5 mg/75 µl)	Effective against bacterial and fungal strain.	[85]
		<i>Staphylococcus aureus</i> , <i>Micrococcus luteus</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus subtilis</i> and <i>Escherichia coli</i> ; <i>Candida albicans</i> and <i>Aspergillus niger</i>	5 mg, 10 mg, 15 mg, 20 mg/75 µl and ciprofloxacin (5 mg/75 µl), Nystatin(5 µg/ml)	Methanolic extract is effective against all microbial while water extract is moderate against all bacterial and fungal strain.	[87]
Methanol and water	[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl] 3-methoxy-4-[3-phenylprop-2-en-1-yl]oxybenzoate dihydrate. (4)				
Anticancer Cancer					
Methanol and water	[6-[[1a,1b,2,5a,6,6a)-6-hydroxy-1a-(hydroxymethyl)-2,5a,6,6a-tetrahydro-1bHoxireno[5,6]cyclopenta[1,3-c]pyran-2-yl]oxy]-3,4,5-trihydroxyoxan-2-yl] methyl-3 phenylprop-2-enoate. (1)	MCF-7 cell	5µg, 25µg, 50µg, 75µg, 100µg/ml and Paclitaxel (5µg/ml)	Inhibit tumor invasion and migration.	[86]
Antidiabetic Activity					
Water	[3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl] 3-methoxy-4-[3-phenylprop-2-en-1-yl]oxybenzoate dihydrate. (5)	Rat	100 mg/kg, PkE, 200 mg/kg and glibenclamide (10mg/kg)	Increase the insulin mediated translocation and expression of glucose transporter-4.	[87]
Hyaluronidase inhibitory activity					
Methanol	Picrorhizaoside-D (11) Picrorhizaoside-E (12)	Mast cell	0µg/ml; 3µg/ml; 10 µg/ml; 30 µg/ml 100 µg/ml and disodium cromoglycate, ketotifenumarate (0,3,10, 30,100µg/ml).	Shown similar or stronger activity than the antiallergic medicines.	[110]
<i>P. scrophulariiflora</i>					
Immunomodulatory activity					
Methanol	scrocaffeside A	Rat	0 µg/ml, 5 µg/ml, 25 µg/ml, 125 µg/ml and Con A(50µg/ml)	shown immunoenhancement effects on immune system	[111]

fat and high-sugar diet for 8 w to establish non-alcoholic steatohepatitis showed antifatty liver effect. Its mechanism appeared to be the regulation of lipid metabolism and reduction of insulin resistance, through inhibition of oxidative stress and inflammation^[112-123].

Most of the biological activities have been studied extensively on *P. kurroa* as compared to *P. scrophulariiflora* as per the available literature, however, few biological activities, which have been

reported were antimicrobial, high metabolic potential, immunomodulation, and therapeutic effect. There is an urgent need to explore the full potential of *P. scrophulariiflora*'s biological activities.

Future scope:

Traditionally rhizomes of *P. kurroa* have been used to cure a liver disorder, fever, lowering sugar level, cancer, asthma, jaundice, malnutrition, urinary

TABLE 4. BIOLOGICAL ACTIVITIES OF PLANT EXTRACTS

Solvent used	Strains/test animals	Doses and control	Result	Reference
<i>Picrorhiza kurroa</i>				
Antimicrobial Activity				
Ethanol; methanol	<i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus subtilis</i> and <i>E.coli</i> ; <i>Candida albicans</i> , <i>Candida tropicalis</i> , <i>Trichophyton rubrum</i> , <i>Aspergillus niger</i> and <i>Penicillium marneffi</i>	10, 25, 50, 100 mg/ml and ciprofloxacin (5 µg/ml), Nystatin (5 µg/ml)	Effective against all the selected bacterial and fungal strain.	[86,101]
Anticancer activity				
Methanol and water	MDA-MB-435S, Hep3B and PC-3 cells	15.625, 31.25, 62.5 and 125 µg/ml and XTT (0.6 mg/ml)	To induce apoptosis	[101]
Ethanol: Water (mother extract) further fractionated using n-hexane, dichloromethane, butanol, acetone, methanol and water	MCF-7, MDA-MB 231 and HeLa, SiHa cell lines	3.9, 7.8, 15.6, 31.2, 62.5, 125, 250, and 500 µg/ml and MTT (5mg/ml)	Dichloromethane fraction of <i>P. kurroas</i> hown effective anticancer activity	[102]
Antidiabetic activity				
Water and ethanol	Rat	100 and 200 mg/kg and glibenclamide 10mg/kg	Helps in b-cell regeneration with enhanced insulin production and antihyperglycemic effects.	[90]
Antiasthmatic Activity				
Ethanol	Guinea pig	25mg/kg And salbutamol (0.2mg/kg)	Reduce the inhibition of histamine and acetylcholine.	[106]
Antianemic activity				
Ethanol	Rat	100 mg/kg/day and 200 mg/kg/day and phenylhydrazine (40mg/kg)	Increase the concentration of haemoglobin and increase the number of red blood cells	[107]
Antioxidant activity				
Methanol and water	Rat	0.5 ml and EDTA	Methanolic extract is less efficient as compared to water in the scavenging of the radicals.	[101]
Methanol	Single cell of hepatocytes	0.7ml/kg and without <i>P. kurroa</i> extract (0.7ml/kg)	Elevated the malondialdehyde levels, decreased superoxide dismutase levels, glutathione levels and oxidative stress.	[108]
Antiarthrits activity				
Water and Ethanol	Rat	2 mL/kg and Indomethacin (3 mg/kg)	Normalize redox status of synovium, suppression of pro-inflammatory cytokines, angiogenesis.	[119]
Antitrypanosomal Activity				

Methanol	Rat	250 mg/kg and 1% gum acacia orally (2 mL/kg)	Reduction and killing of trypanosomes.	[109]
Hepatoprotective activity				
Ethanol and Water	Rat	400 to 1500 mg/day and without extract	Significant restoration of liver enzyme level.	[92]
<i>P. scrophulariiflora</i> Therapeutic effect				
Water	Rat	20, 40 and 80 mg/kg/d and CMC (0.5%)	Significantly decreased total cholesterol and triglycerides assay in liver tissue and increased the Superoxide dismutase activity.	[112]

disorder, leukoderma, snake bite, piles, and rheumatic diseases. To verify the credibility of traditional uses, biological activities of rhizomes are yet to be tested in the areas of leukoderma, snake bite, and hemorrhoids. Similarly, the seeds and flowers of *P. kurroa* have been used traditionally to treat liver and upper respiratory tract disorders, fever, dyspepsia, chronic diarrhea, and blood purification but there is not much supporting data has been generated. Several traditional uses have been recorded to cure a liver disorder, choleric, joint pain, asthma, jaundice, arthritis, and bacterial infection from the stems of *P. scrophulariiflora* but no scientific study has been carried out in this regard either.

In *P. kurroa* there are 29 chemical constituents (6-10,13-26,36-45), which have not been tested for any biological activities. Similarly, 9 chemical constituents (27-35) are isolated from *P. scrophulariiflora* but none were assessed for biological activities. Some biological activities of the genus *Picrorhiza* that differ from traditional uses are antianemic activity, antitrypanosomal activity, and high metabolic activity. Only some biological activities have buttressed traditional uses such as antimicrobial, hepatoprotective activity, antiasthmatic activity, arthritis activity, anticancer activity, antidiabetic activity.

Phytochemical studies on the genus *picrorhiza* have revealed at least 53 compounds in the last decade while the biological activities have been carried out on 8 compounds. These studies reveal antimicrobial, anticancer, antidiabetic, hyaluronidase inhibitory, immunomodulatory activities. On the basis of literature, it could be concluded that the rhizomes possessed a good profile of chemical constituents and biological activities. However more studies are required to explore about *P. scrophulariiflora*.

Acknowledgments:

Authors would like to thank Maharishi Markandashwar Educational Trust, Sadopur-Ambala (Haryana), India for their support in all respect.

Conflict of Interest:

The authors declare that they have read the policy and guidelines of the journal and there are no conflicts of interest.

REFERENCES

1. Sumner J. The Natural History of Medicinal Plants, Timber Pres: London; 2000. p. 69.
2. Heinrich M, Barnes J, Gibbons S, Williamson EM. Fundamentals of Pharmacognosy and Phytotherapy. UK: Churchill Livingstone, Elsevier Science Ltd; 2004. p. 336.
3. Scurlock J. Sourcebook for Ancient Mesopotamian Medicine. Atlanta: SBL Press; 2014. p.786.
4. Samant SS, Dhar U, Palni LMS. Medicinal plants of IndianHimalaya: Diversity, distribution and potential value. Nainital: Gyano-daya Prakashan; 1998. p. 163.
5. Naser SH. Islamic Science-An illustrated study. Westerham. England: Westerham Press; 1976. p. 15.
6. Ahmad M, Khan MA, Marwat SK. Useful medicinal flora enlisted in Holy Quran and Ahadith. Am Eurasian J Agric Env Sci 2009;5(1):126-40.
7. Yari KH, Kazemi E, Yarani R, Tajehmiri A. Islamic Bioethics for Fetus Abortion in Iran. Am J Sci Res 2011;18:118-21.
8. Craig SR. Windhorses and Dharma Warriors: The Religious, Historical, and Cultural Significance of Horse Protection Rituals in Mustang, Nepal. In: Olsean SL, Littauer MA, Ingrid R, editors. Horses and Humans: The Evolution of Human-equine Relationships BAR International Series: Oxford; 2006. p. 339-54.
9. Mark B. Duke's Handbook of Medicinal Plants of the Bible, HerbalGram. Austin, United States: American Botanical Council; 2010. p. 69.
10. Sharma S. Realms of Ayurveda. New Delhi: Arnold- Heineman; 1979.
11. Bhowmik D, Kumar KPS, Srivastava S, Paswan S, Sankar A, Dutta D. Traditional Indian herbs punarnava and its medicinal importance. J Pharmacogn Phytochem 2012;1:52-7.

12. Tucakov J. Healing with plants – Phytotherapy. Beograd: Culture; 1971. pp. 180-90.
13. Jain SK, De Phillips RA. Medicinal Plants of India. vol. 2. Algonac (MI): Reference Publications; 1991. p. 558.
14. Varier PS. Indian Medicinal Plants. A Compendium of 500 Species. vol. 4. Hyderabad. India: Orient Longman; 1995. pp. 269-72.
15. Singh DK, Hajra PK. Floristic diversity. In: G.S. Gujral, V. Sharma V. Changing perspective of biodiversity status in the Himalaya, British Council Division, Wildlife Youth Services. New Delhi: British High Commission Publication; 1996. p. 23-38.
16. Sharma PV. Classical Uses of Medicinal Plants. Varanasi: Chaukhamba Visvabharati; 1996. p. 69-70.
17. Fakim-Gurib A. Medicinal plants: traditions of yesterday and drugs of tomorrow. Mol Aspect Med 2006;1-93.
18. Simbo DJ. An ethnobotanical survey of medicinal plants in Babungo. Northwest Region: Cameroon. J Ethnobiol Ethnomed 2010;6:1-7.
19. Chintamunnee V, Mahomoodally MF. Herbal medicine commonly used against infectious diseases in the tropical island of Mauritius. J Herb Med 2012;2:113-25.
20. Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. Front Pharmacol 2014;4:177
21. Joshi RK, Satya IP, Setzer WN. Himalayan aromatic medicinal plants: a review of their ethnopharmacology, volatile phytochemistry and biological activities. Medicines 2016;3(1):1-55.
22. Masood M, Arshad M, Qureshi R, Sabir S, Amjad MS, Qureshi H, Tahir Z. *Picrorhiza kurroa*: an ethnopharmacologically important plant species of the Himalayan region. Pure Appl Biol 2001;4:407-17.
23. Schippmann U, Leaman DJ, Cunningham AB. Impact of cultivation and gathering of medicinal plants on biodiversity, in: Global Trends and Issues, Biodiversity and the ecosystem approach in agriculture, forestry and fisheries. Rome: Commission on Genetic Resources for Food and Agriculture; 2002. p. 1-21.
24. Ramakrishnappa K. Impact of cultivation and gathering of medicinal plants on Biodiversity: Case studies from India, in: Biodiversity and the Ecosystem Approach in Agriculture, Forestry and Fisheries. 2002. Available from: <http://www.wwf.org.uk/filelibrary/pdf/medplantsandcons.pdf>
25. Wang L, Ren J, Zhe G. Current status and future direction of Chinese herbal medicine. Trends Pharmacol Sci 2002;23:348-48.
26. Hamilton AC. Medicinal plants, conservation and livelihoods. Biodiversity Conservation 2004;13:1477.
27. Sasidharan S, Chen Y, Saravanan D, Sundram KM, Yoga L. Extraction, isolation and characterization of bioactive compounds from plants extracts. Afr J Tradit Complement Altern Med 2011;8:1-10.
28. Petrovska BB. Historical review of medicinal plants usage. Pharmacog Rev 2012;6:1-5.
29. Mondal TK, Bantawa P, Sarkar B, Ghosh P, Chand PK. Cellular differentiation, regeneration, and secondary metabolite production in medicinal *Picrorhiza* spp. Plant Cell Tissues Organ Cult 2013;112(2):143-58.
30. Sah JN, Varshney VK. Chemical constituents of *Picrorhiza* genus: a Review. Am J Essential Oil 2013;1(2):22.
31. Kumar N, Kumar T, Sharma SK. Phytopharmacological review on genus *Picrorhiza*. Int J Universal Pharm Bio Sci 2013;2(4):334-47.
32. Katoch M, Fazli IS, Suri KA, Ahuja A, Qazi GN. Effect of altitude on picroside content in core collections of *Picrorhiza kurroa* from the north western Himalayas. J Nat Med 2011;65:578-82.
33. Royle JF. Illustrations of the botany and other branches of the natural history of the Himalayan mountains and of the flora of Cashmere. New Delhi: Volume plates, Reprint, Today and Tomorrow's Printers and Publishers; 970. p. 71.
34. Coventry BO. Wild flowers of Kashmir Vol. 2. Dehradun: Reprint 1984. Bishen Singh Mahendra Pal Singh; 1927. p. 89-90.
35. Barrie FR, Burdet HM, Chaloner WG, Demoulin V. International code of botanical nomenclature, adopted by the fifteenth international botanical congress, Yokohana, Koeltz Scientific Books: Konigstein; 1994. p. 52-3.
36. Subedi B, Shretha R. Plant Profile. Himalayan. Bioresour 1999;3:14-15.
37. Polunin O, Stainton A. Flowers of the Himalayas. 6th ed. Delhi: Oxford University Press; 1990. p. 295.
38. Gaddipati JP, Madhavan S, Sidhu GS. Picroliv – a natural product protects cells and regulates the gene expression during hypoxia/reoxygenation. Mol Cell Biochem 1999;194:271-81.
39. Prajapati ND, Purohit SS, Sharma AK, Kumar T. A handbook of medicinal plants: A complete source book. Jodhpur, India: Agrobios; 2003. p. 396-97.
40. Singh GB, Sarang B, Singh S, Khajuria A, Sharma ML, Gupta BD, et al. Antiinflammatory activity of the iridoids kutkin,- picroside-1 and kutkoside from *Picrorhiza kurroa*. Phytotherapy Res 2006;7:402-07.
41. Stuppner H, Wagner H. New cucurbitacin glycosides from *Picrorhiza kurroa*. Planta Med 1989;55:559-63.
42. Dorsch W, Stuppner H, Wagner H, Grapp M, Demoulin S, Ring J. Antiasthmatic effect of *Picrorhiza kurroa*: Androsin prevents Allergen and PFA- induced Branchial obstruction in Guinea pigs. Int Arch Allergy Immunol 1991;95:128-33.
43. Chettri N, Sharma E, Lama SD. Non-timber forest produces utilization, distribution and status in a trekking corridor of Sikkim. India. Lyonia 2005;8:94-108.
44. Bantawa P, Ghosh SK, Moitra S, Ghosh P, Mondal TK. Status and conservation threats of *Picrorhiza scrophulariiflora* Pennell. (Scrophulariaceae): An endangered high valued medicinal plant of the Indo-China Himalayan Region. Bioremed Biodiver Bioavail 2009;3:15-22.
45. Pennell FW. The Scrophulariaceae of the Western Himalayas. Reprint 1997. Dehradun, India: Bishen Singh Mahendra Pal Singh; 1943. p. 63-66.
46. Bantawa P, Saha-Roy O, Ghosh SK, Mondal TK. *In vitro* regeneration of an endangered medicinal plant *Picrorhiza scrophulariiflora*. Biol Plant 2011;55:169-72.
47. World Health Organization. African Traditional Medicine: African Technical Report Series No 1, Congo Brazaville: WHO Regional Office for Africa; 1976.
48. Maroyi A. An ethnopharmacological survey of medicinal plants used by the people in Nhema communal area. Zimbabwe. J Ethnopharmacol 2012;136(2):347-54.
49. Mahwasane ST, Middleton N, Boaduo N. An ethnobotanical survey of indigenous knowledge on medicinal plants used by traditional healers of the Lwamondo area: Limpopo province South Africa. S Afr J Bot 2013;88:69-75.
50. Kashyap SR. Samhita or Vriddha Jivakiya Tantra. Trans IGM Shastri. Mumbai, India: Bombay Sastu Sahitya; 1970.

51. Gulabkunverba S. Charak Sutra 26, Charak Samhita. Jamnagar: India: Ayurvedic Society; 1949. p. 124-41.
52. Kirtikar KR, Basu BD. Indian Medicinal Plants. In: Harathi K, Naidu CV, editors. Influence of Ethylene Inhibitor Silver Nitrate on Direct Shoot Regeneration from in Vitro Raised Shoot Tip Explants of *Sphaeranthus indicus* Linn.—An Important Antijaundice Medicinal Plant. Vol. III. Allahabad: Lalit Mohan Basu; 1935. p. 1825-26.
53. Chopra RN, Chopra IC, Handa KL, Kapur LD. Chopra's Indigenous Drugs of India, second edition. Calcutta: Dhur and Sons; 1981. pp. 181-83.
54. Chaturvedi GN, Singh RH. Jaundice of infections hepatitis and its treatment with an indigenous drug *Picrorhiza kurroa*. J Res Ind Med 1966;1:1-5.
55. Krishnamurthy A. The Wealth of India. New Delhi: Council of Industrial and Scientific Research; 1969. p. 49.
56. Weinges K, Kloss P, Henkels WD Natural products from medicinal plants, XVII, Picroside –II, a new 6- Vanilloyl-catapol from *Picrorhiza kurroa* Royle and Benth. Justus Liebigs Ann Chem 1972;759:173-82.
57. Nadkarni AK. "Indian Meteria Medica". Mumbai: Popular Parkashan Pvt. Ltd.; 1976. p. 953-954.
58. Pandey GS. Bhavaprakasa Nighantoo (Indian Materia Medica). Varanasi: India: Chaukhamba Sanskrit Sansthan; 1979. p. 70–71.
59. Dey AC. Indian Medicinal Plants Used in Ayurvedic Preparations. Dehradun, India: Bhishen Singh Mahendra Pal Singh; 1980. p. 11.
60. Langer JG, Gupta OP, Atal CK. Clinical trials *Picrorhiza kurroa* as immunomodulator. Indian J Pharmacol 1981;13:98–99.
61. Yegnarayan R, Dange SV, Vaidya SD, Balwani M. Study of *Picrorhiza kurroa* (PK300) in case of bronchial asthma. Bombay Hosp J 1982;24:81-84.
62. Jayaweera DMA. Medicinal Plants Used in Ceylon, Part 5. Colombo. Sri Lanka: The National Science Council of Sri Lanka; 1982. p. 76.
63. Mogre K, Vora KK, Sheth UK. Effect of *Picrorhiza kurroa* and *Eclipta alba* on Na⁺K⁺ATPase in hepatic injury by hepatotoxic agent. Indian J Pharmacol 1982;13:253-59.
64. Atal CK, Sharma ML, Kaul A, Kuajuria A. Immunomodulating agents of plant origin. I: Preliminary screening. J Ethnopharmacol 1986;18(2):133-41.
65. Ansari RA, Aswal BS, Chander R, Dhawan BN, Garg NK. Hepatoprotective activity of kutkin, the iridoid glycoside mixture of *Picrorhiza kurroa*. Indian J Med Res 1988;87:401-04.
66. Zhang H, Zheng Z. Handbook of Chinese Traditional Medicine Resources. Beijing: Science Press; 1994. p. 1149.
67. Rajaram D. A preliminary clinical trial of *Picrorhiza kurroa* in bronchial asthma. Bombay Hosp J 1996;18:66–69.
68. Luper S. A review of plants used in the treatment of liver disease: part 1. Altern Med Rev 1998;3(6):410-421.
69. Gupta PP. Picroliv Hepatoprotectant, immunomodulator. Drug Fut 2001;26:25-31
70. Anonymous. Ayurvedic interventions for diabetes mellitus: a systematic review. Altern Med Rev 2001;6:319-21.
71. Liu J, Liu BL, Zhang JQ, Zhang N. Hepatoprotective and choleric action of *Picrorhiza scrophulariae* flora Pennell. Chin J New Drug 2002;11:459-61.
72. Zhang Y, DeWitt DL, David S, Muraleedharan NG. Cyclooxygenase-2 enzyme inhibitory triterpenoids from *Picrorhiza kurroa* seeds. Life Sci J 2005;77(25):3222–30.
73. Kala CP. Medicinal plants of the high altitude cold desert in India: Diversity, distribution and traditional uses. Int Bio Sci Manage; 2006;2:43-56.
74. Girish C, Pradhan SC. Drug development for liver diseases: Focus on picroliv, ellagic acid and curcumin. Fundam Clin Pharmacol 2008;22(6):623-32.
75. Bhandari P, Kumar N, Singh B, Ahuja PS. Online HPLC-DPPH method for antioxidant activity of *Picrorhiza kurroa* Royle ex Benth and characterisation of kutkoside by ultra-performance LC electrospray ionization quadrupole time of flight mass spectroscopy. Indian J Exp Biol 2009;48(3):323-28.
76. Phondani PC, Maikhuri RK, Kala CP. Ethnoveterinary uses of medicinal plants among traditional herbal healers in alankanda catchment of Uttarakhand, India. Afr J Tradit Complem 2010;7:195-206.
77. Bisht VK, Kandari LS, Negi JS, Bhandari AK, Sundriyal RC. Traditional use of medicinal plants in district Chamoli, Uttarakhand, India. J Med Plant Res 2013;7(15):918-29.
78. Ghisalberty EL. Biological and pharmacological activity of naturally occurring iridoids and secoiridoids. Phytomedicine 1998;5(2):147-63.
79. State Pharmacopeia. Commission of P. R. China, Pharmacopeia of the P. R. China. Beijing: People's Health Publishing House; 1995. p. 204.
80. Zou ALC, Zhu TF, Gan SC, Wang DC, Zhang Y, Deng XM. Two new secoiridoid glycosides Zhongguo Yiyao Gongye Zazhi from the roots *Picrorhiza scrophulariiflora*. Chin Chem Lett 2008;19:1224-27.
81. Bhattacharjee S, Bhattacharya S, Jana S, Bhagel DS. A Review on Medicinally Important Species of *Picrorhiza*. Intl J Pharm Res Biosci 2013;2(4):1-16.
82. C P C Pharmacopoeia of the People's Republic of China, Beijing: Chemical Industry Press; 2005. p. 167.
83. Goyal BR, Agrawal BB, Goyal RK, Mehta AA. Pharmacological classification of herbal anti-asthmatics. Orient Pharm Exp Med 2007;7:11-2580.
84. Lohani H, Gwari G, Andola HC, Chauhan N. High performance liquid chromatographic determination of Picroside I & II in *Picrorhiza kurroa* Royle ex Benth: A critically endangered plant of Himalaya. Med Plants 2010;3:165-67.
85. Mohammed R, Mohammed U, Yamgar S, Gadgoli C, Salunkhe D. Preliminary screening and antimicrobial activity of *Picrorhiza kurroa* royle ethanolic extracts. Int J Pharm Sci Rev Res 2012;14(1):73-76.
86. Rathee D, Thanki M, Bhuvra S, Anandjiwala S, Agrawal R. Iridoid glycosides-Kutkin, Picroside I and Kutkoside from *Picrorrhiza kurroa* Benth inhibits the invasion and migration of MCF-7 breast cancer cells through the down regulation of matrix metalloproteinases. Arab J Chem 2013;6(1):49-58.
87. Husain GM, Rai R, Rai G, Singh HB, Thakur AK, Kumar V. Potential mechanism of anti-diabetic activity of *Picrorhiza kurroa*. Tang 2014;4(4):1-5.
88. Kumar V, Sood H, Chauhan RS. Optimization of a preparative RP-HPLC method for isolation and purification of picrosides in *Picrorhiza kurroa*. J Plant Biochem Biot 2016;25(2):208-14.
89. Sultan AJ, Pervaiz Q. Phytochemical studies for quantitative estimation of iridoid glycosides in *Picrorhiza kurroa* Royle. Bot Stud 2016;57(7):4-6.
90. Kumar S, Patial V, Soni S, Sharma S, Pratap K, Kumar D, Padwad Y. *Picrorhiza kurroa* Enhances b-Cell Mass Proliferation and Insulin Secretion in Streptozotocin Evoked b-Cell Damage in Rat. Front Pharmacol 2017;8(537):1-15.

91. Ali M, Sultana S, Mir Rasool S. Chemical Constituents from the Roots of *Picrorhiza kurroa* Royle Ex Benth. *Int J Pharm Pharm Sci* 2017;9(3):25-35.
92. Navya K, Kumar GP, Chandrasekhar Y, Kumar A. Evaluation of potassium dichromate (K₂Cr₂O₇)-induced liver oxidative stress and ameliorative effect of *Picrorhiza kurroa* extract in Wistar Albino Rats. *Biol Trace Elem Res* 2018;184:(1)154-64.
93. Win NN, Kodamab T, Zar Khine LW, Winc YY, Ngwec H, Ikuro A, Hiroyuki M. Bis-iridoid and iridoid glycosides: Viral protein R inhibitors from *Picrorhiza kurroa* collected in Myanmar. *Fitoterapia* 2019;134:101-07.
94. Sacchettini JC, Poulter CD. Creating isoprenoid diversity. *Science* 1997;277:1788-1789.
95. Xie Z, Hu H, Kong D, Yang P Two new compounds from *Picrorhiza scrophulariiflora*. *Chin J Pharma* 2007;38(3):221-26.
96. Zou X, Liao X, Ding LS, Peng SL. Phenyl and phenylethyl glycosides from *Picrorhiza scrophulariiflora*. *J Asian Nat Prod Res* 2007;9(3-5):443-48.
97. Huang KY, He L, Qu Y, Gao HY, Deng XM, Wu LJ Isolation and identification of chemical constituents from roots of *Picrorhiza scrophulariiflora* Pennell. *Shenyang Yaoke Daxue Xuebao* 2009;26(2):112-15.
98. Zhu TF, Jiang K, Yan ZH, Hu JJ, Zou LC, Wang DC, Deng XM. A new secoiridoid glycoside from roots of *Picrorhiza scrophulariiflora*. *Chin Herbal Med* 2013;5(3):237-39.
99. Xiong K, Juc Z, Tong Z, Wangc Z, Han H. Metabolic profiles and pharmacokinetics of picroside I in rats by liquid chromatography combined with electrospray ionization tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 2018;1095:157-65.
100. Kumar V, Sharma N, Shitiz K, Singh TR, Tan C, Sood H, Chauhan RS. An insight into conflux of metabolic traffic leading to picroside-I biosynthesis by tracking molecular time course changes in a medicinal herb, *Picrorhiza kurroa*. *Plant Cell Tissue Organ Cult* 2015;123:435-41.
101. Rajkumar V, Gunjan G, Kumar A; Antioxidant and anti-neoplastic activities of *Picrorhiza kurroa* extracts. *Food Chem Toxicol* 2011;49(2):363-69.
102. Mallick MN, Singh M, Parveen R, Khan W, Ahmad S, Najm MZ, Husain SA. HPTLC Analysis of Bioactivity Guided Anticancer Enriched Fraction of Hydroalcoholic Extract of *Picrorhiza kurroa*. *BioMed Res Int* 2015;1-18.
103. Hussain A, Shadma W, Maksood A, Ansari SH. Protective effects of *Picrorhiza kurroa* on cyclophosphamide-induced immunosuppression in mice. *Pharmacog Res* 2013;5(1):30-35.
104. Nagarathna PKM, Reena K, Reddy S, Wesley J. Review on immunomodulation and immunomodulatory activity of some herbal plants. *Int J Pharmaceut Sci Rev Res* 2013;22(1):223-30.
105. Shubha KS, Sumana K, Lakshmidivi L. Antifungal activity of *Solanum xantocarpum* Sch and Wend and *Picrorhiza kurroa* Royle ex Benth against some clinical dermatophytes. *Int J Curr Microbiol Appl Sci* 2016;5(2):236-44.
106. Sehgal R, Chauhan A, Gilhotra UK, Gilhotra A. *In vitro* and *in vivo* evaluation of antiasthmatic activity of *Picrorhizz kurroa* Plant. *Int J Pharm Sci* 2013.4(9):3440-43.
107. Turaskar A, More S, Sheikh R, Gadhpayle J, Bhongade SL. Inhibitory potential of *Picrorhiza kurroa* royle ex. benth extracts on phenylhydrazine induced reticulocytosis in rats. *Asian J of Pharm Clin Res* 2013;6(2):215-16.
108. Deshpande N, Das RK, Manohar M, Das V, Kandi S, Ramana KV. Antioxidant effects of *Picrorhiza kurroa* rhizome extracts in alcoholic cirrhosis of liver. *Am J Pharmacol Sci* 2015;3(5):49-51.
109. Shaba P, Pandey NN, Sharma OP, Rao J, Singh RK. *In vitro* antitypanosomal evaluation of *Picrorhiza karroa* rhizome. *Int Res J Pharm* 2012;3(7):205-08.
110. Morikawaa T, Nakanishia Y, Inouea H, Mansea Y, Matsuuraa H, Hamasakia S, *et al.* Acylated iridoid glycosides with hyaluronidase inhibitory activity from the rhizomes of *Picrorhiza kurroa* Royle ex Benth. *Phytochemistry* 2020;169:112-85.
111. An N, Wang D, Zhu T, Zeng S, Cao Y, Cui J, *et al.* Effects of scrocaffeside A from *Picrorhiza Scrophulariiflora* on immunocyte function *in vitro*. *Immunopharmacol Immunotoxicol* 2009;31(3):451-58.
112. Xu X, Wang W, Zhao Z, Xi W, Yu B, Hao C, *et al.* Effects of total iridoid glycosides of *Picrorhiza scrophulariiflora* against non-alcoholic steatohepatitis rats induced by high-fat and high-sugar diet through regulation of lipid metabolism. *Chin Herb Med* 2020;42:1-6.
113. Firdous J, Bharathi V, Muhamad N, Zaini F. Evaluation of anti-microbial activity in *Picrorhiza kurroa* plant extract using thin-layer chromatography and FTIR. *Int J Pharm Tech* 2016;8(3):15717-722.
114. Zou L, Yan Z, Zhu T, Xiang H, Wang D. Two new phenylethanoid glycosides and a new secoiridoid glycoside from the roots of *Picrorhiza scrophulariiflora*. *Chin Chem Lett* 2010;21:1103-06.
115. Zhu T, Chen J, Yang Y, Sun Q, Hu L, Yan ZH, Li P. Secoiridoid glycosides from the roots of *Picrorhiza scrophulariiflora*. *Chem Nat Compd* 2018;54(4):23.
116. Wang H, Zhao W, Choomuenwai V, Andrews K, Quinn R, Feng Y. Chemical investigation of an antimalarial Chinese medicinal herb *Picrorhiza scrophulariiflora*. *Bioorg Med Chem Lett* 2013;23:5915-18.
117. Usman M, Surekha Y, Chhaya G, Devendra S. Preliminary screening and antimicrobial activity of *Picrorhiza kurroa* Royle ethanol extracts. *Int J Pharm Sci Rev Res* 2012;14:73-76.
118. Rathee D, Rathee P, Rathee S, Rathee D. Phytochemical screening and antimicrobial activity of *Picrorrhiza kurroa*, an Indian traditional plant used to treat chronic diarrhea. *Arab J Chem* 2016;9:1307-13
119. Kumar R, GuptaYK, Singh S, Arunraja S. *Picrorhiza kurroa* inhibits experimental arthritis through inhibition of Pro-inflammatory Cytokines, Angiogenesis and MMP. *Phytother Res* 2016;30:112-19.
120. Krupashree K, Kumar K, Rachitha P, Jayashree GV, Khanum F. Chemical composition, antioxidant and macromolecule damage protective effects of *Picrorhiza kurroa* Royle ex Benth. *S Afr J Bot* 2014;94:249-54.
121. Cao X, Qiao J, Wang L, Ye X, Zheng L, Jiang N, Mo W. Screening of glycoside isomers in *P. scrophulariiflora* using ionic liquid-based ultrasonic-assisted extraction and ultra-performance liquid chromatography/electrospray ionization quadruple time-of-flight tandem mass spectrometry. *Rapid Commun Mass Spectrom* 2012;26:740-48.

122. Chen Z, Wu J, Shen W, Liu P, Cao Y, Lu Y. Counter-current chromatographic method for preparative scale isolation of picrosides from traditional Chinese medicine *Picrorhiza scrophulariiflora*. *J Separation Sci* 2011;34(15):1910-16.
123. Tiwari SS, Pandey MM, Srivastava S, Rawat AKS. TLC densitometric quantification of picrosides (picroside-I and picroside-II) in *Picrorhiza kurroa* and its substitute *Picrorhiza scrophulariiflora* and their antioxidant studies. *Biomed Chromatogr* 2012;26:61-68.
-