

# Indian Journal of Pharmaceutical Sciences

Scientific Publication of the Indian Pharmaceutical Association

Indexed in Ind MED, EMBASE/Excerpta Medica, International Pharmaceutical Abstracts, Chemical Abstracts.

Volume 70

Number 1

January-February 2008

## CONTENTS

### REVIEW ARTICLES

#### A Decision Tree for Rapid Quality Assurance and Control of Rifampicin-Containing Oral Dosage Forms for Global Distribution for Tuberculosis Treatment

Y. ASHOKRAJ, SHRUTIDEVI AGRAWAL AND R. PANCHAGNULA 1-4  
SWATI RAWAT, SUDHA VENGURLEKAR, B. RAKESH, S. JAIN, G. SRIKARTI 5-10

### RESEARCH PAPERS

#### *In vivo* Evaluation of Single Dose Tetanus Toxoid Vaccine Formulation with Chitosan Microspheres

R. MANIVANNAN, S. A. DHANARAJ, Y. UDAYA BHASKARA RAO, A. BALASUBRAMANIAM, N. L. GOWRISHANKAR, N. JAWAHAR AND S. JUBIE 11-15

#### Ionic Cross-linked Chitosan Beads for Extended Release of Ciprofloxacin: *In vitro* Characterization

A. SRINATHA, J. K. PANDIT AND S. SINGH 16-21

#### Design and Optimization of Diclofenac Sodium Controlled Release Solid Dispersions by Response Surface Methodology

H. N. SHIVAKUMAR, B. G. DESAI AND G. DESHMUKH 22-30

#### Evaluation of Free Radical Scavenging Activity of an Ayurvedic Formulation, *Panchvalkala*

SHEETAL ANANDJIWALA, M. S. BAGUL, M. PARABIA AND M. RAJANI 31-35

#### Validation of Different Methods of Preparation of *Adhatoda vasica* Leaf Juice by Quantification of Total Alkaloids and Vasicine

S. SONI, SHEETAL ANANDJIWALA, G. PATEL AND M. RAJANI 36-42

#### Formulation and Characterization of Mucoadhesive Buccal Films of Glipizide

MONA SEMALTY, A. SEMALTY AND G. KUMAR 43-48

#### Synthesis, Antimicrobial and Anti-inflammatory Activity of 2,5-Disubstituted-1,3,4-oxadiazoles

G. NAGALAKSHMI 49-55

#### Ascorbic Acid Inhibits Development of Tolerance and Dependence to Opiates in Mice: Possible Glutamatergic or Dopaminergic Modulation

S. K. KULKARNI, C. DESHPANDE AND A. DHIR 56-60

#### Design and *In Vitro* Characterization of Buccoadhesive Drug Delivery System of Insulin

J. SAHNI, S. RAJ, F. J. AHMAD AND R. K. KHAR 61-65

#### Development and Evaluation of a Chloramphenicol Hypertonic Ophthalmic Solution

A. V. JITHAN, C. KRISHNA MOHAN, AND M. VIMALADEVI 66-70

#### Optimization of Fast Dissolving Etoricoxib Tablets Prepared by Sublimation Technique

D. M. PATEL AND M. M. PATEL 71-76

#### Furosemide-loaded Alginate Microspheres Prepared by Ionic Cross-linking Technique: Morphology and Release Characteristics

M. K. DAS AND P. C. SENAPATI 77-84

### SHORT COMMUNICATIONS

#### Isolation of Liver Aldehyde Oxidase Containing Fractions from Different Animals and Determination of Kinetic Parameters for Benzaldehyde

R. S. KADAM AND K. R. IYER 85-88

#### Microwave-Induced Synthesis of Schiff Bases of Aminothiazolyl Bromocoumarins as Antibacterials

K. N. VENUGOPALA AND B. S. JAYASHREE 88-91

#### *In vitro* Antiviral Activity of some Novel Isatin Derivatives against HCV and SARS-CoV Viruses

P. SELVAM, N. MURGESH, M. CHANDRAMOHAN, E. DE CLERCQ, E. KEYAERTS, L. VIJGEN, P. MAES, J. NEYTS AND M. V. RANST 91-94

#### Physicochemical and Pharmacokinetic Parameters in Drug Selection and Loading for Transdermal Drug Delivery

N. S. CHANDRASHEKAR AND R. H. SHOBHA RANI 94-96

#### HPLC Estimation of berberine in *Tinospora cordifolia* and *Tinospora sinensis*

G. V. SRINIVASAN, K. P. UNNIKRISHNAN, A. B. REMA SHREE AND INDIRA BALACHANDRAN 96-99

#### Parenteral Formulation of Zopiclone

P. V. SWAMY, P. SUSHMA, G. CHIRAG, K. PRASAD, M. YOUNUS ALI AND S. A. RAJU 99-102

#### Simultaneous Spectrophotometric Determination of Lansoprazole and Domperidone in Capsule Dosage Form

A. P. SHERJE, A. V. KASTURE, K. N. GUJAR AND P. G. YEOLE 102-105

#### Novel 2-Pyrazoline Derivatives as Potential Antibacterial and Antifungal Agents

SUVARNA KINI AND A. M. GANDHI 105-108

#### Spectrophotometric Estimation of Ethamsylate and Mefenamic Acid from a Binary Mixture by Dual Wavelength and Simultaneous Equation Methods

ANJU GOYAL AND I. SINGHVI 108-111

#### Novel Colon Targeted Drug Delivery System Using Natural Polymers

V. RAVI, T. M. PRAMOD KUMAR AND SIDDARAMAIAH 111-113

#### Effect of Some Clinically Used Proteolytic Enzymes on Inflammation in Rats

A. H. M. VISWANATHA SWAMY AND P. A. PATIL 114-117

#### Synthesis and Pharmacological Evaluation of (6-Substituted 4-Oxo-4H-chromene-3 yl) methyl N-substituted Aminoacetates

ASMITA GAJBHIYE, V. MALLAREDDY AND G. ACHAIHAH 118-120

#### Development and *In Vitro* Evaluation of Buccoadhesive Tablets of Metoprolol Tartrate

P. D. NAKHAT, A. A. KONDAWAR, L. G. RATHI AND P. G. YEOLE 121-124

#### RP-HPLC Estimation of Venlafaxine Hydrochloride in Tablet Dosage Forms

S. L. BALDANIA, K. K. BHATT, R. S. MEHTA, D. A. SHAH AND TEJAL R. GANDHI 124-128

#### Simultaneous Estimation of Esomeprazole and Domperidone by UV Spectrophotometric Method

S. LAKSHMANA PRABU, A. SHIRWAIKAR, ANNIE SHIRWAIKAR, C. DINESH KUMAR, A. JOSEPH AND R. KUMAR 128-131

#### *In Vitro* Anthelmintic Activity of *Baliospermum montanum* Muell. Arg roots

R. G. MALI AND R. R. WADEKAR 131-133

#### REFEREES FOR INDIAN JOURNAL OF PHARMCEUTICAL SCIENCES DURING 2006 & 2007

134-134

# *In vivo* Evaluation of Single Dose Tetanus Toxoid Vaccine Formulation with Chitosan Microspheres

R. MANIVANNAN\*<sup>1</sup>, S. A. DHANARAJ, Y. UDAYA BHASKARA RAO<sup>2</sup>, A. BALASUBRAMANIAM<sup>1</sup>, N. L. GOWRISHANKAR<sup>1</sup>, N. JAWAHAR<sup>1</sup> AND S. JUBIE<sup>1</sup>

J. S. S. Centre for Research and Post Graduate Studies, J. S. S. College of Pharmacy, Ootacamund - 643 001,

<sup>1</sup>Swamy Vivekanandha College of Pharmacy, Elayampalayam (Thiruchengode) - 637 205, <sup>2</sup>Pasteur Institute of India, Coonoor - 643 103, India

Manivannan, *et al.*: *In vivo* Evaluation of Single Dose Tetanus Toxoid Vaccine

Chitosan adsorbed microspheres containing tetanus toxoid were prepared in the size range of 10  $\mu\text{m}$  to 75  $\mu\text{m}$ , by emulsion-cross linking technique at different speeds of agitation. The amount of tetanus toxoid incorporated into chitosan microspheres were estimated by limes flocculation test and *in vivo* evaluation of tetanus toxoid adsorbed chitosan microspheres were determined by toxin neutralization method using albino mice. The results of *in vivo* release for the batches of 10  $\mu\text{m}$  and 25  $\mu\text{m}$  correlates with the results of *in vitro* in which both the batches passes the limit of IP standard (4 Lf) where as, for the batches of 50  $\mu\text{m}$  and 75  $\mu\text{m}$ , the *in vitro* release of tetanus toxoid was 2 Lf. But our *in vivo* studies for the batches of 50  $\mu\text{m}$  and 75  $\mu\text{m}$  fail to pass the limit stated in IP. The release of tetanus toxoid from the chitosan microspheres was found to be sustained for the period of 6 months.

**Key words:** Tetanus toxoid, microspheres, chitosan, tetanus toxin, tetanus antitoxin

Controlled delivery of bioactive macromolecules has become increasingly important. Vaccines are one of the important classes of molecules that could potentially benefit from controlled release because many immunization schedules require administration of several spaced antigen doses. In particular the conversion of multiple-dose vaccines into single-dose vaccines may represent an important advancement, which should lead to improved vaccination coverage, as well as to a reduction in vaccination costs<sup>1</sup>.

Many vaccines against pathogenic infections such as tetanus, diphtheria, pertussis, poliomyelitis and hepatitis B require more than one dose for complete immunization of a child. The multidose schedule of immunization increases the cost of immunization several fold, as well as having the disadvantages of inconvenience and incomplete subject compliance. To overcome these shortcomings, technologies have been developed to formulate single contact point delivery systems for vaccines requiring multiple injections<sup>2</sup>. This project is to develop a controlled release tetanus toxoid vaccine formulation, which could replace the

two or three doses, required for primary immunization series.

The initial priority was to develop a single-dose tetanus vaccine, as one of the most urgent needs of the Expanded Programme on Immunization is to control neonatal tetanus. According to World Health Organization (1999) statistics, each year, more than 4 50 000 infants die from neonatal tetanus (tetanus in the first month of life), and nearly 40 000 mothers die from tetanus infection acquired during delivery. So, the prevention of neonatal tetanus would greatly benefit from the development of vaccine requiring only one injection in pregnant women or women childbearing age. However, this technology if successful, will also be applied to other vaccines, namely DPT and hepatitis B and possibly inactivated polio vaccine. For this work, tetanus toxoid has been selected because of its characteristics and availability and appears as a suitable antigen to begin alternative approaches to the existing vaccines.

Tetanus Toxoid is a protein product prepared from tetanus toxin from the growth of a highly toxicogenic strain of tetanus bacilli, *C. tetani* in a suitable medium. The concentrated and purified tetanus toxoid is adsorbed on a mineral adjuvant

---

\*For correspondence

E-mail: manivannan\_biotech@yahoo.co.in

(aluminium hydroxide/aluminium phosphate) to enhance considerably its immunizing activity<sup>3</sup>.

Chitosan is a high molecular weight, cationic polysaccharide obtained by the deacetylation of chitin<sup>4</sup>, the major compound of exoskeletons of crustaceans. Because of its low production costs, biodegradability, biocompatibility, bioadhesivity, non-toxic<sup>5,6</sup> in nature and recent FDA approval, the pharmaceutical and food applications of chitosan have increased remarkably over recent years.

## MATERIALS AND METHODS

Tetanus toxoid (700 L<sub>f</sub>/ml), tetanus toxin, tetanus antitoxin and tetanus antitoxin-national reference was obtained from Pasteur Institute of India, Coonoor. Chitosan was a gift sample from Central Institute of Fisheries Technology, Cochin. Linseed oil, toluene, glutaraldehyde (AR Grade), were purchased from S. D. Fine Chemicals Ltd, Mumbai. All the other chemicals used in the present study were of AR Grade. For conducting *in vivo* experiments, animals were procured from the central Animal House of the Institute. All animal experimental protocols were approved by the Institutional Animals Ethics Committee.

### Preparation of chitosan-gel microspheres<sup>7</sup>:

Chitosan-gel microspheres were prepared by emulsion-crosslinking technique in which glutaraldehyde was used as the cross-linking agent. The solutions of 1.5% chitosan and 2.0% chitosan were prepared in aqueous acetic acid (3%) containing sodium chloride (2%) and this was stirred at 3000 rpm to form a gel. A dispersion phase was prepared by mixing 50 ml of toluene and 5 ml of span-80 using stainless steel remistirrer at 1000 rpm speed for 10 min. To this 4 ml of chitosan gel with 1 ml of 0.01 N hydrochloric acid was added and the stirring was continued, then glutaraldehyde saturated toluene (5 ml) was added drop wise. Finally 4% glycine (2 ml) was added to cap any free aldehyde groups. The microspheres were collected and washed in acetone and toluene with centrifugation and they were dried at room temperature and stored in desiccator.

### Loading of tetanus toxoid:

The tetanus toxoid was incorporated into chitosan gel microspheres by adsorption method. In the preparation of tetanus toxoid loaded chitosan microspheres,

**TABLE 1: DIFFERENT BATCHES OF FORMULATION WERE USED FOR *IN VIVO* STUDIES**

Different batches of formulation
Tetanus toxoid adsorbed onto 1.5% chitosan microspheres (10 µm)
Tetanus toxoid adsorbed onto 1.5% chitosan microspheres (25 µm)
Tetanus toxoid adsorbed onto 2.0% chitosan microspheres (50 µm)
Tetanus toxoid adsorbed onto 2.0% chitosan microspheres (75 µm)
Mixed batches (1.5% and 2.0%)
Pasteur Institute Vaccine (Control)

Six different batches of formulation using 1.5% (10 µm and 25 µm) and 2.0% (50 µm and 75 µm) Chitosan microspheres were used for *in vivo* studies

400 mg of the pooled 1.5% empty microspheres were mixed with 40 ml of sterile water and stored for 24 h, to achieve proper swelling. The swelled microspheres were separated by centrifugation at 7000 rpm then washed with acetone. These microspheres were immersed in about 4 ml of plain tetanus toxoid (700 L<sub>f</sub>/ml) and kept at 37° for 20 h adsorption. The microspheres were centrifuged and unadsorbed supernatant tetanus toxoid (0.75 ml) was removed. Then the tetanus toxoid adsorbed 1.5% chitosan microspheres were washed with methanol and acetone and they were taken for further evaluation<sup>8</sup>. Three other batches of microspheres were prepared by the above mentioned method with 2% chitosan. The Table 1 shows the different batches of tetanus toxoid adsorbed chitosan microspheres were used for *in vivo* studies.

The amount of tetanus toxoid incorporated into chitosan microspheres were estimated by limes flocculation (L<sub>f</sub>) test. Limes flocculation means, when the concentration of toxin, or toxoid, is kept constant and the concentration of the antitoxin varied in mixtures of constant volume, the mixture flocculating first is that which contains the most nearly equivalent quantities of toxin, or toxoid and antitoxin<sup>9</sup>.

### *In vivo* evaluation of microspheres:

The suggested method in the Indian Pharmacopoeia is to determine the potency of tetanus vaccine is antibody induction method (toxin neutralization method). The various steps are involved in the antibody induction method, immunization of guinea pigs, determination of Lp/10 dose, collection of immune sera from guinea pigs, titration of standard antitoxin (control titration) and titration of immune guinea pig sera at Lp/10 level.

### Immunization of guinea pigs<sup>10</sup>:

The guinea pigs for test and control group were selected. The animal ranged from weights 250-350 g and quarantined for 1 w before use with regular diet.

The tetanus toxoid adsorbed chitosan microspheres were dispersed in aqueous vehicle (0.5% sorbitol, 0.1% carboxymethylcellulose and 0.002% Tween-80) to get a final concentration of 0.5 Lf/ml. These six samples were injected subcutaneously into six groups of guinea pigs.

#### Collection of immune sera from guinea pigs<sup>11</sup>:

Bleeding of the immunized guinea pigs were done by cardiac puncture under general anaesthesia (with anesthetic ether) using 20 gauge needle. The blood samples were collected in sterile glass tubes and kept in a slanting position to allow the serum to ooze out of the clotted blood. Then these tubes containing clotted blood and serum were kept at 37° for 1 h and stored at 4° in for 24 h. The serum was separated by centrifugation at 2500 rpm for 15 min and the serum of each guinea pig accordingly labeled and stored at -20° before performing neutralization test.

#### Determination of *Limes paralyticum*/10 (Lp/10) dose of the test toxin:

The *Limes paralyticum*/10 dose of the test toxin is the smallest quantity of the toxin, which mixed with 0.1 unit of the standard preparation and injected subcutaneously into mice causes tetanic paralysis within 4 d. For the determination Lp/10 dose, the mixtures were prepared by adding 2.0 ml of diluted solution of the tetanus antitoxin-national reference preparation into each of five graded volumes of diluted test toxin correspondingly to 2.32, 2.40, 2.48, 2.56, and 2.64 ml. Their volumes were made up to 5.0 ml with peptone water, so that 0.5 ml of each mixture contained 0.1 unit of tetanus antitoxin. Mixtures were incubated at room temperature protected from 15-60 min. After incubation, the mice were inoculated with 0.5 ml of mixture per mouse and the injected mice were observed for 4 d.

#### Titration of standard antitoxin (control titration):

Control titration was carried out along with the neutralization test to check the validity of the Lp/10 test. From the solution of diluted standard antitoxin, five different concentrations were prepared by adding correspondingly 1.6, 1.8, 2.0, 2.2 and 2.4 ml of diluted standard antitoxin. To this 2.0 ml of diluted toxin was added and the volume was made upto 5.0 ml with peptone water, so that 0.5 ml of each mixture contained 0.0060 ml of toxin. The mixtures were incubated at room temperature, protected from light for 15-60 min and then inoculated into each mouse.

#### Titration of immune guinea pig sera at Lp/10 level:

The neutralization test was carried out on pooled sera to obtain the end point (the international unit per ml), the pooled sera was diluted further accordingly such that the end point value was determined for unitage in different intervals of bleeding.

## RESULTS AND DISCUSSION

From our studies, it was observed that the size of microspheres increased with the decrease in the speed of agitation. The amount of drug loaded was 70-84% by limes flocculation test. The biodegradable microspheres made up of chitosan polymer having good spherical geometry, adsorbed onto tetanus toxoid (plain) was used for protracted pulsed release of the incorporated toxoid to reduce multiple immunization regimes. The size of the microspheres was found to be ranging between 10-75 µm.

From our studies, 0.006 ml (Table 2) of standard toxin dose per mouse was the smallest quantity of the toxin which when mixed with 0.1 unit of

**TABLE 2: DETERMINATION OF Lp/10 (LIMES PARALYTICUM/10) DOSE OF THE TEST TOXIN**

Toxin dose (ml)	Antitoxin dose (TNR) IU	Mixtures table				Observation			
		Dil. toxin (ml)	Dil. antitoxin (ml)	Peptone water (ml)	Total volume (ml)	1 <sup>st</sup> day	2 <sup>nd</sup> day	3 <sup>rd</sup> day	4 <sup>th</sup> day
0.0058	0.1	2.32	2.0	0.68	5.0	✓✓✓	✓✓✓	✓✓✓	t tt tt,
						✓✓✓	✓✓✓	✓✓✓	t t tt
0.006	0.1	2.40	2.0	0.60	5.0	✓✓✓	✓tt	✓tt	tt tt ttt,
						✓✓✓	✓✓✓	✓✓✓	t ttt ttt
0.0062	0.1	2.48	2.0	0.52	5.0	✓✓✓	t t tt	t t tt	D D D
						✓✓✓	tt tt ttt	tt tt ttt	
0.0064	0.1	2.56	2.0	0.44	5.0	✓✓✓	ttt ttt ttt	ttt ttt ttt	
						✓✓t	ttt ttt tt	ttt ttt tt	
0.0066	0.1	2.64	2.0	0.36	5.0	✓✓✓	D ttt ttt	D ttt ttt	
						✓tt	ttt ttt ttt	ttt ttt ttt	

✓ - Normal mouse, t- tetanic paralysis of one limb, tt- tetanic paralysis of two limb, ttt- tetanic paralysis of three limb, D- death due to tetanus

**TABLE 3: MAXIMUM POTENCY UNITAGE ESTIMATED FOR DIFFERENT VACCINE FORMULATIONS AT DIFFERENT LEVEL**

Different vaccine formulations	Unitage (IU/ml)			
	2 <sup>nd</sup> week after booster dose	6 <sup>th</sup> week after booster dose	9 <sup>th</sup> week after booster dose	12 <sup>th</sup> week after booster dose
1.5% TT adsorbed chitosan batch (10 µm)	>0.5	2.0	2.0	1.0
1.5% TT adsorbed chitosan batch (25 µm)	>0.5	2.0	2.0	1.5
2.0% TT adsorbed chitosan batch (50 µm)	>0.5	0.5	<0.5	<0.5
2.0% TT adsorbed chitosan batch (75 µm)	>0.5	0.5	<0.5	<0.5
Mixed batches (10 + 25 + 50 + 75 µm)	>0.5	1.0	1.0	1.0
Control batch (PIIC)	>0.5	2.0	4.0	4.0

TT is tetanus toxoid, PIIC - Pasteur Institute of India, Coonoor

the standard antitoxin preparation caused tetanic paralysis with in 4 d and this was used as the Lp/10 dose. After ascertaining the level of Lp/10 value of tetanus toxin, the standard titration was done using the same quantity of tetanus toxin (0.006 ml) and different quantities of standard antitoxin (0.1 IU/ml). The Table 3 shows the tetanus toxoid adsorbed onto different size (10 µm, 25 µm, 50 µm, 75 µm and mixed batches) of chitosan microspheres have elicited specific antibodies at different level.

The 10 µm microspheres of 1.5% chitosan have elicited a specific antibody level of 0.5 IU/ml to 2.0 IU/ml in the sera for 2<sup>nd</sup> and 6<sup>th</sup> w after the booster dose, respectively. The antibody level of 2.0 IU/ml remained the same in 9<sup>th</sup> w serum sample. Although the immune response elicited by 1.5% tetanus toxoid adsorbed chitosan (10 µm) batch has decreased to 1.0 IU/ml after the 12<sup>th</sup> w of booster dose, but it has passed the Indian Pharmacopoeia standard <sup>9</sup>.

The 25 µm microspheres of 1.5% chitosan also elicited the same response as 10 µm microspheres, but the antibody level after 12 w was did not go below 1.5 IU/ml as in the case of 10 µm batch. The 50 µm and 75 µm of 2.0% chitosan microspheres exhibited comparatively a lower antibody response, having the maximum antibody level of 0.5 IU/ml after 6 w. For these microsphere batches after 12 w the antibody level was reduced below 0.5 IU/ml. This may be due to the delay or obstructed release of antigen from the microsphere matrix.

The mixed batches (1.5% and 2.0%) containing all four different sizes of microspheres adsorbed with tetanus toxoid have elicited a lower antibody response when compared to the individual batches (10 µm and 25 µm). In this batch, the antibody response remained constant at 1 IU/ml after 6 w.

From these observations it was ascertained that 10 µm, 25 µm, and mixed batches of chitosan adsorbed were able to produce the antibodies to the required level mentioned in official monograph, but when the antibody level produced was compared with Pasteur Institute Vaccine (control) the antibody level was 50% less. Since for Pasteur Institute Vaccine (control), the antibody level was 2-4 IU/ml. From these results indicated that chitosan microspheres are capable of inducing required antibody titre, it is necessary to study the factors which can improve the microspheres system for eliciting the antibody level equivalent to Pasteur Institute Vaccine (control) and also for a longer antibody induction period.

The present study suggested that the maintenance of antibody level has to be estimated over a period of 6 mo so as to confirm the pulsatile releasing mechanism of chitosan microspheres for the enhancement of immunogenicity and for proving the efficiency of vaccine delivery system.

## ACKNOWLEDGEMENTS

The authors wish to thank Central Institute of Fisheries Technology, Cochin for providing the generous gift sample of Chitosan. Authors also thank Pasteur Institute of India, Coonoor for the facilities provided and for the encouragement.

## REFERENCES

1. Alonso MJ, Gupta RK, Caroline Min, Siber GR, Langer R. Biodegradable microspheres as controlled release tetanus toxoid delivery systems. *Vaccine* 1994;12:299.
2. Diwan M, Misra A, Khar RK, Talwar, GP. Long-term high immune response to diphtheria toxoid in rodents with diphtheria toxoid conjugated to dextran as a single contact point delivery system. *Vaccine* 1997;15:1867.
3. Christensen PE. Third Inter Conf Tetanus, Sao Paulo; Brazil: 17-22, 1970;PAHO/WHO Scientific publication, No.253, 1972. p. 36.
4. Illum L. Chitosan and its use as a pharmaceutical excipient. *Pharm Res* 1978;15:1326.

5. Felt O, Buri P Gurny R. Chitosan: A unique polysaccharide for drug delivery. *Drug Develop Ind Pharm* 1998;24:979.
6. Paul W, Sharma CP. Chitosan: A drug carrier for the 21<sup>st</sup> century; A review. *STP Pharma Sci* 2000;10:5.
7. Yuichi O, Shiratani M, Kobayashi H, Ouchi T. Release behavior of 5-fluorouracil from chitosan-gel microspheres immobilizing 5-fluorouracil coated with polysaccharides and their cell specific cytotoxicity. *Pure Appl Chem* 1994;31:629.
8. Jayakrishnan A, Jameelaa SR, Misra A. Cross-linked chitosan microspheres for prolonged delivery of macromolecular drugs. *J Biomater Sci Polymer Edn* 1994;6:621.
9. Indian Pharmacopoeia Vol II, Delhi: Controller of Publications; 1996. p. A-33.
10. Calcagno ML, Manghi MA. A possible explanation for the discrepancy between ELISA and neutralising antibodies to tetanus toxin. *Vaccine* 2000;18:2698.
11. Gupta RK, Alroy J, Alonso MJ, Langer R, Siber GR. Chronic local tissue reactions, long term immunogenicity and immunologic priming of mice and guineapigs to tetanus toxoid encapsulated in biodegradable polymer microspheres composed of polylactide-co-glycolide polymers. *Vaccine* 1997;15:1716.

**Accepted 6 January 2008**

**Revised 10 September 2007**

**Received 24 July 2006**

**Indian J. Pharm. Sci., 2008, 70 (1): 11-15**